

PABRON KIDS COLD GRANULES

- ◆ PABRON KIDS COLD GRANULES is a cold medication for children that has steady effect on cough, nasal and throat symptoms, and fever.
- ◆ The particles are designed for kids -i.e., flavored with strawberries that kids love for easy administration, and does not contain caffeine that disturbs sleep.
- ◆ Acute cough, fever, and runny nose should be relieved at any cost. Treat with PABRON KIDS COLD GRANULES at an early stage, keep the body warm, and have a good rest.

Indication

To alleviate cold symptoms (cough, runny nose, fever, sore throat, sneezing, sputum, stuffy nose, chills (feeling cold due to fever), headache, joint pain, and muscle pain)

Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

7 to 10 years: 1 packet per dose, 3 times daily

3 to 6 years: 2/3 packet per dose, 3 times daily

1 to 2 years: 1/2 packet per dose, 3 times daily

Under 1 year: Do not take

Precaution

- Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.
- Infants under 2 years old should seek medical care from a physician first. Use the drug only in an unavoidable case.

Ingredient

In 1 packet (0.96g)

Tipepidine Hibenstate 12.5mg

Guaifenesin 25mg

Chlorpheniramine Maleate 1.2mg

Acetaminophen 150mg

Excipients

Anhydrous silicic acid, corn starch, sugar, D-Mannitol, aspartame (L-phenylalanine compound), acesulfame K, sucralose, hydroxypropylcellulose, erythrosine, magnesium stearate, flavoring, vanillin, ethyl vanillin

Precautions

Although the drug is intended for children, the document describes the general precaution for cold medicine.

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur)

- This product should not be taken in the following persons
 - Patients who have had an allergic symptom to this drug or its ingredients.
 - Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.
- Do not use any of the following medications while taking this drug
 - Other cold medicines, antipyretics/analgesics, sedatives, expectorants, and internal medicines, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.)
- After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness may occur)
- Do not drink alcohol before and after taking this drug
- Do not take this medicine for a long time

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration
 - Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms.

Hyperthermia, urination difficulty

Persons diagnosed as having the following.

Heart disease, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Skin: rash/redness, itching

Gastrointestinal system: nausea/vomiting, and loss of appetite

Neuropsychiatric system: dizziness

Urinary system: urination difficulty

Other: excessive temperature decrease

- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomuocutaneous syndrome (stevens-johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Drug-induced hypersensitivity syndrome: Symptoms such as widespread redness of the skin, generalised rash, fever, malaise, and swelling of lymph nodes (neck, armpit, groin, etc.) may occur.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.

Interstitial pneumonia: Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance.

Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.

Aplastic anemia: symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.

Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Dry mouth, drowsiness

- When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you

Precautions for storage and handling

- Avoid direct sunlight, and store the product in a cool place with little humidity.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.
- Do not take the product past the expiration date.

31

[Disclaimer on Multilingual Product Information]

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.