

# BIOFERMIN®

## ANTIDIARRHOIC

Class 2 OTC drug

Antidiarrheals

● Relieves abdominal pain due to diarrhea "Scopolia Extract" and "Peony root extract" alleviate abdominal pain due to diarrhea. ● Protects gastrointestinal organs damaged by diarrhea. "Berberine tannate" and "Geranium herb dry extract" protect damaged intestinal mucosa and relieve diarrhea. ● Gently relieves intestinal conditions "Bifidobacterium bifidum powder" improves stool condition (watery stool), and gently relieves intestinal conditions. ● Easy to take, convenient to carry around Pale yellowish brown to yellowish brown tablets have little bitterness or odor, and are easy to take. They are packed in PTP sheets convenient to carry around.

### Indication

diarrhea accompanied by abdominal pain, Diarrhea, diarrhea due to indigestion, food poisoning, water poisoning, vomiting and purging, loose bowels and soft stools.

### Dosage and administration

Take the following dose with cold or lukewarm water after a meal.

15 years old or over: 3 tablets per dose, 3 times daily

11 to 14 years old : 2 tablets per dose, 3 times daily

Under 11 years old : Do not take

### Precautions of Dosage and Administration

The take of the medicine in children should be supervised by a parent.

Strictly follow the recommended dosage and administration.

How to take out the tablets

• Press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets.

(If taking the tablets with the blister pack, it will lead to unexpected accidents such as the sharp edge sticking into the esophageal mucosa.)

### Ingredient and amount

In 9 tablets (daily dosage for persons  $\geq$  15 years old)

Berberine Tannate 300mg

Geranium herb dry extract 420mg (extracted from about 4200 mg of the crude medicine)

Scopolia Extract 33mg

Peony Root extract 125mg (extracted from 500 mg of the crude medicine)

Bifidobacterium bifidum powder 30mg

### Excipients

Corn starch ,Maltose Syrup powder,Carmellose Calcium,Talc,Magnesium Stearate

### Precautions relating to ingredient

The medicine contains herbal medicine (medicinal leaves, roots, barks, etc.). The color and taste of the tablets may differ slightly, but this will not affect the efficacy.

### Precautions

#### To be avoided

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

● This medicine should not be taken together with any of the following medicines:

Pain relief/antispasmodic medicine for the stomach, other gastrointestinal medicine containing scopolia extract, medicine for motion sickness

● After taking this medicine, do not drive a car or operate machinery (symptoms such as blurred vision and abnormal brightness may occur).

● Nursing women should either not take this medicine or stop nursing while taking this medicine. (the medicine may be transferred to the breast milk and accelerate the pulse of the infant. )

### Consultation

● The following persons should consult a physician, pharmacist, or registered salesperson before taking this medicine.

Patients undergoing medical treatment from a physician

Those who have diarrhea with fever, bloody stool or persistent mucous stool.

Those who have acute and severe diarrhea or diarrhea with accompanying symptoms such as abdominal pain/abdominal distension/nausea. (The disease may worsen by taking this medicine to stop diarrhea. )

Pregnant women or women suspected of being pregnant.

The elderly

Patients who have experienced allergic symptoms associated with medicines, etc.

Persons with the following symptoms:

urination difficulty

Persons diagnosed as having the following

heart disease, Glaucoma, Persons on oral medicines containing parasympathetic blocker (belladonna total alkaloid, isopropamide iodide, scopolia extract, etc.)

● If the following symptoms are observed after taking this medicine, these may be adverse reactions, so immediately discontinue the use of this medicine, and show package insert to your physician, pharmacist, or registered salesperson for a consultation.

Skin: rash/redness, itching,

Neuropsychiatric system: headache

Urinary system: urination difficulty

Other: Facial hot flashes, abnormal brightness

● The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and show package insert to your physician, pharmacist, or registered salesperson for a consultation.

Dry mouth, blurred vision

● When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take package insert with you.

### Other precautions

● Breast milk may be difficult to produce.

### Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with low humidity.

Store the product beyond the reach of children.

Do not transfer the medicine to other containers. (It may lead to misuse or quality deterioration)

Do not take the product past the expiration date.

Enter the date when the inner pouch (aluminum pouch) was opened in the "date of opening" section on the box.

After opening the inner pouch (aluminum pouch), consume the content as soon as possible hopefully within 6 months after opening the product to ensure product quality.

11

#### 【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product (or quasi medicine) approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.