

Make sure to read this instruction sheet before use.
Additionally, store this sheet so that it can be referenced when needed.

Before scratching and leaving eczema wounds on your hands that you don't want people to notice



MEDIROID PVA CREAM

CLINILABO

Class 2 OTC drug

Brand name: CLINILABO MEDIROID PVA CREAM

<Therapeutic agent for eczema and dermatitis>

- ◆ **CLINILABO MEDIROID PVA CREAM** is a medicine containing a medium-grade steroid component that is effective against ailments such as eczema of the hand that you don't want to people to notice.
- ◆ The four active ingredients relieve painful eczema and dermatitis, one of which is prednisolone valerate acetate, an antedrug steroid that works on the affected area and has low activity in the body.
- ◆ The cream is smooth and non-sticky, for ailments such as eczema of the hands.



Precautions for Use



Things You Must Not Do



(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects occurring)

- 1 Do not use on the following areas:
 - (1) Varicella (chicken pox), fungal infections such as athlete's foot or ringworm, or areas affected by suppuration.
 - (2) Around the eyes, mucous membranes, etc.
- 2 Do not use extensively on the face
- 3 Do not use for long periods of time



Things To Consult On



- 1 The following individuals should consult with their doctor, pharmacist, or a registered seller before using this medicine
 - (1) Persons receiving treatment from a doctor.
 - (2) Persons who are pregnant or who think they may be pregnant.
 - (3) Persons with a history of allergic reactions caused by medicines, etc.
 - (4) Persons whose skin is extensively affected.
 - (5) Persons with damp skin or severe sores.

- 2 If any of the following symptoms appear after you have used this medicine, you may be experiencing side effects. As such, please discontinue use immediately and consult with your doctor, pharmacist or registered seller, bringing a copy of this instruction sheet with you

Relevant Area	Symptoms
Skin	Rash/redness, itching, contact dermatitis, feeling of dryness, irritation, heat sensation, stinging sensation
Skin (affected area)	Types of tinea such as athlete's foot and ringworm, acne, suppuration symptoms, and persistent irritation

- 3 If symptoms do not improve after 5 to 6 days of using this medicine, discontinue use and consult with your doctor, pharmacist, or registered seller with a copy of this instruction sheet

Efficacy

Eczema, dermatitis, heat rash, contact dermatitis, itching, insect bites, hives

Usage/Dosage

Take an appropriate amount and apply it to the affected area several times a day.


Note:

- (1) Strictly follow the prescribed usage and dosage.
- (2) If this medicine is being used by a child, ensure it is used under the guidance and supervision of a parent or guardian.
- (3) Avoid contact with eyes. In case of contact, rinse with cold or lukewarm water immediately. If you experience severe symptoms, see an eye specialist.
- (4) Use only for topical application.
- (5) Do not cover the area where this medicine is applied with anything that has poor breathability, such as cling film.

Handling of the aluminum tube

- Squeeze from the folded part of the tube (end) towards the tip.

(①→②)



- The tube may tear and the medicine may spill out.

Do not bend.



Do not fold.



Components

Per 1 g

Components	Amount	Action
Prednisolone valerate acetate	1.5 mg	Relieves the source of inflammation and suppresses eczema, itching, and dermatitis.
Crotamiton	50 mg	Quickly soothes itching.
Tocopherol acetate	5 mg	Promotes blood circulation and aids in recovery of the affected area.
Isopropylmethylphenol	1 mg	Sterilizes the affected area to prevent bacterial infection.

Additives: Polysorbate 60, polyoxyethylene hydrogenated castor oil, propylene glycol, parabens, stearyl alcohol, cetanol, carboxyvinyl polymer, diethyl sebacate, triethanolamine

Precautions for Storage and Handling

- (1) Ensure the cap is tightened securely and store the medicine in a cool place that is out of direct sunlight.
- (2) Store this product out of reach of children.
- (3) Do not transfer to another container. (This may result in misuse or a change in quality)
- (4) Do not use this product beyond its expiration date. Use this medicine as soon as possible after opening, even if still within the expiration date. (In order to ensure quality.)



Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.