DERMARIN POWDER SPRAY DX

Ringworm Remedies

Indication

Tinea pedis, tinea cruris, tinea corporis

Dosage and administration

Spray an appropriate dose onto the lesion once daily.

Comply with the prescribed dosage and administration instructions.

Do not leave the affected lesion and surrounding area soiled when using the medicine.

Attention should be paid so that the medicine will not come into contact with eyes. In case the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water, and immediately seek an ophthalmologist for medical treatment.

When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it. This medication is only for external use.

Do not touch the eyes and other mucous membranes with fingers stained with the drug.

Shake well before use. Spray on the affected site at a distance of approximately 10 cm. Confirm the position of the spray nozzle to avoid spraying the drug on the face, particularly toward the eyes, and to prevent inhalation.

Do not spray the drug to the same site for more than 1 second. (The drug cools the site significantly, and excessive coldness may cause frostbite.)

ingredient and amount

In 90 g Terbinafine Hydrochloride 135mg Lidocaine 270mg Glycyrrhetinic Acid 67.5mg I-Menthol 135mg

Excipients

Sorbitan sesquioleate ,Octyldodecyl myristate,Anhydrous ethanol,Magnesium carbonate ,Titanium oxide,Magnesium aluminometasilicate ,Corn starch ,LPG,Isopentane

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.) • This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Do not use in the following sites.

Eyes and periorbital area, mucosa (for example, oral cavity, nasal cavity, vagina, etc.), scrotum, vulvar, etc. eczema

Lesions with severe moistening, erosion, fissures, and trauma.

Consultation

• The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use. Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

infants and small children

Patients who have experienced allergic symptoms associated with drugs, etc.

The lesion is on the face or an extensive area.

Patients with purulent hemorrhoidal lesions.

Those whose symptoms are not differentiated between "eczema" or "athlete's foot, jock itch, ringworm". (when symptoms such as itchiness/erosion are present on the scrotum, they are often attributed to other causes such as eczema.)

• If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and take this container to your physician, pharmacist, or registered sales person for a consultation. Skin: eruption, irritation, feeling hot, scale/exfoliation (peeled skin such as dandruff and scurf), sores, dryness/feeling of tension, skin fissure, pain, pigmentation, rash/redness*, itching*, swelling*, urticaria* (*: These symptoms may develop in the entire body.)

If symptoms do not resolve after using the drug for about 2 weeks or if symptoms worsen owing to the use of the drug, stop taking this drug, and take this container to your physician, pharmacist, or registered salesperson for a consultation.

Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not use the product past the expiration date (to assure quality).

Do not place the drug close to any source of flame.

Be careful in handling the drug because it may degrade the paint of the floor and furniture when it touches.

If you turn the can sideways when the remaining drug becomes small, the drug may not come out smoothly.

[Disclaimer on Multilingual OTC Product Information]

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.