Make sure to read this instruction sheet before use. Additionally, store this sheet so that it can be referenced when needed.

For blurred vision and eye strain

Class 3 OTC drug **IRIS 40** <Eve Drops>

- In daily life, we overuse our eyes, causing symptoms such as trouble seeing things up close, blurred vision causing difficulty reading texts, and a tendency toward eye strain.



IRIS 40 activates the metabolism of the eyes, returns eye function toward its normal state, and is effective against symptoms such as blurred vision and eye strain. Precautions for Use Things To Consult On The following individuals should consult with

- their doctor, pharmacist, or a registered seller before using this medicine.
- (1) Persons who are currently being treated by a doctor.
- (2) Persons who have had allergic symptoms due to medication in the past.
- (3) Persons with the following symptoms:
- Intense eye pain (4) Persons who have received any of the following diagnoses: Glaucoma

If any of the following symptoms appear after you have used this medicine, you may be experiencing side effects. As such, please discontinue use immediately and consult with your doctor, pharmacist or registered seller, bringing a copy of this instruction sheet with you.

Relevant Area	Symptoms	
Skin	Rash, redness, itching	
Eyes	Red eye, itching, swelling	

- In case of the following, discontinue use and consult with your doctor, pharmacist, or registered seller, bringing a copy of this instruction sheet with you.
 - (1) Your blurred vision does not improve.
 - (2) Your symptoms do not improve after using the product for about 2 weeks.

Efficacy

Blurred vision (e.g. with excessive eye discharge), eye strain, itchy eyes, conjunctival hyperemia, eye inflammation caused by UV rays or other light rays (e.g. snow blindness), prevention of eye diseases (e.g. after swimming, dust and sweat in the eyes), blepharitis (inflammation of the eyelids), and discomfort when wearing hard contact lenses

For these occasions:



When you have eye strain caused by driving



For blurred vision and eye strain during computer use, etc.



For eye strain after watching TV

Usage/Dosage

Apply to eyes 3 to 6 times daily, 2 to 3 drops each time.

Note:

- (1) Strictly observe the prescribed dosage and administration.
- (2) If this medicine is being used by a child, ensure it is used under the guidance and supervision of a parent or guardian.
- (3) Do not let the tip of the container touch your eyelids or eyelashes. Do not use if the solution is cloudy.
- (4) Do not use while wearing soft contact lenses.
- (5) Use only for application in the eyes.

	Ingredients		
Components	Amount	Action	
Neostigmine methylsulfate	0.002%	Improves your eyes' weakened ability to adjust their focus.	
D-α-tocopherol acetate (natural vitamin E)	0.03%	Antioxidant and blood circulation promoting effects improve blood flow and aid in the supply of nutrition.	
Flavin adenine Sodium dinucleotide (active vitamin B2)	0.05%	Promotes optimal eye function by stimulating metabolism.	
Taurine	1.0%	Nourishes the eyes, promotes metabolism, and is effective for eyes strain.	
Chondroitin sulfate sodium	0.1%	Protects the eye by preventing the cornea from drying out.	
Chlorpheniramine maleate	0.02%	Antihistamine effect to relieve itching in eyes.	

Excipients: L-menthol, Dryobalanops aromatica resin, benzalkonium chloride, chlorobutanol, sodium edetate, polysorbate 80, polyoxyethylene hydrogenated castor oil, tonicity agent, boric acid, citric acid, sodium citrate

Precautions for Storage and Handling

- Ensure the cap is tightened securely and store the medicine in a cool place that is out of direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not replace the medicine into a different container. (This may result in misuse or a change in quality)
- (4) Do not share eye drops with anyone else.
- (5) Do not place eye drops in high-temperature locations, such as on the dashboard of your car. (The container may become deformed or the quality of the solution may deteriorate.)
- (6) If the drug spills and stains your clothes while applying to the eye, immediately wash with water. (The yellow color of this drug is due to the ingredient vitamin B₂.)
- (7) Do not use this product beyond its expiration date. Use this medicine as soon as possible after opening, even if still within the expiration date. (In order to ensure quality.)
- (8) The ingredients may crystallize on the tip of the container or inside the cap depending on the storage conditions. If this occurs, wipe the product gently with clean gauze or similar material before using.



Precautions and Correct Technique for Eye Drop Application

Precautions

 Avoid contact between the tip of the container and your eyelids or eyelashes. Eye discharge or microorganisms may contaminate the solution or cause it to become cloudy.



Applying the Product Correctly

- Wash your hands thoroughly and avoid touching your eyes directly with your fingers.
- Look up slightly and pull your lower eyelid down with your fingers. Apply drops onto the inner side of your eyelid.

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy
 and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your
 information only. It does not warrant that its contents and the product itself conforms to laws and regulations in
 countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified
 or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.