# IRIS NEO <SOFT>

Class 3 OTC drug

**Ophthalmic Medicines** 

◆IRIS NEO <soft> is an eye drop that relieves ocular fatigue and blurry vision due to abuse of the eyes in daily life. ■ "Neostigmine methylsulfate" improves the focus adjustment of the eye, and relieves ocular fatigue. ■ Fresh and soft sensation of use.

# Indication

Eyestrain, prevention of eye troubles (after swimming, or to wash out sweat or dust etc.), foreign-body feeling by contact lenses, blurred vision (eye mucus)

# Dosage and administration

Instill 2 to 3 drops per dose 3 to 6 times daily.

Comply with the prescribed dosage and administration instructions.

When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it. Do not touch the tip of the container with the eyelid or eyelash. Do not use the drug when the solution becomes turbid. Do not use the drug when you are wearing soft contact lenses.

Use this only as an eye drop.

#### ingredient and amount

Neostigimine methylsulfate 0.005% d-a-Tocopherol acetate (Natural vitamin E) 0.05% Pyridoxine Hydrochloride (vitamin B6)(Vitamin B6) 0.1% Taurine (Aminoethyl Sulfonic Acid) 1% Potassium L-aspartate 1%

#### Excipients

I-Menthol, Dryobalanops aromatica resin, Chlorobutanol, Benzalkonium chloride, citric acid, Sodium citrate, Boric acid, pH adjusting agent, Edetate sodium, Polyoxyethylene hydrogenated castor oil, Polysorbate 80, Anhydrous ethanol

#### Precautions

## Consultation

• The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use. Patients undergoing medical treatment from a physician

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

Severe eye pain

Persons diagnosed as having the following

Glaucoma

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation. Skin

rash/redness

itching

Eyes:

congestion of the eye

itching

swelling

• In case of the following, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

When blurred vision does not improved.

When symptoms do not improve even after using the medicine for about 2 weeks

## Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with an air tight closure on.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)

Do not share the drug with other people.

Do not leave the drug under high temperatures, such as on a car dashboard. (It may result in the deformation of the container and the degradation of the drug solution.)

Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).

The crystalized ingredient may accumulate at the tip of the container and inside the cap depending on the storage conditions. Wipe the crystalized ingredient with a clean gauze.

[Disclaimer on Multilingual OTC Product Information]

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.