

◆ IRIS PC effectively relieves ocular fatigue after staring at the monitor of a personal computer and mobile devices for long hours. ■
Soft and cool sensation of use.

Indication

Eyestrain, redness of the conjunctiva, prevention of eye troubles (after swimming, or to wash out sweat or dust etc.) , ophthalmia by ultraviolet rays etc. (snow blindness etc.), blepharitis (running eye), foreign-body feeling by contact lenses, itchy eyes, blurred vision (eye mucus)

Dosage and administration

Instill 2 to 3 drops per dose 3 to 6 times daily.
Comply with the prescribed dosage and administration instructions.
Use for children should be supervised by a parent.
Do not touch the tip of the container with the eyelid or eyelash. Do not use the drug when the solution becomes turbid.
Do not use the drug when you are wearing soft contact lenses.
Use this only as an eye drop.

ingredient and amount

Neostigimine methylsulfate 0.005%
Allantoin 0.3%
Taurine (Aminoethyl Sulfonic Acid) 1.00%
Potassium L-aspartate 0.8%
Chondroitin Sulfate Sodium 0.10%
Pyridoxine Hydrochloride (vitamin B6) 0.10%
d-α-Tocopherol acetate 0.05%

Excipients

l-Menthol,dl-Camphor,Mentha oil ,Chlorobutanol ,Benzalkonium chloride,citric acid,Sodium citrate,Boric acid,pH adjusting agent,Edetate sodium,Hypromellose,Polyoxyethylene hydrogenated castor oil,Polysorbate 80,Anhydrous ethanol

Precautions

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.
 - Patients undergoing medical treatment from a physician
 - Patients who have experienced allergic symptoms associated with drugs, etc.
 - Persons with the following symptoms:
 - Severe eye pain
 - Persons diagnosed as having the following
 - Glaucoma
- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered sales person for a consultation.
 - Skin
 - rash/redness
 - itching
 - Eyes:
 - congestion of the eye
 - itching
 - swelling
- In case of the following, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.
 - When blurred vision does not improved.
 - When symptoms do not improve even after using the medicine for about 2 weeks

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with an air tight closure on.
Store the product beyond the reach of children.
Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)
Do not share the drug with other people.
Do not leave the drug under high temperatures, such as on a car dashboard. (It may result in the deformation of the container and the degradation of the drug solution.)
Do not take the product past the expiration date.
Even within the expiration date, use up as soon as possible once it is opened (to assure the quality).

The crystalized ingredient may accumulate at the tip of the container and inside the cap depending on the storage conditions. Wipe the crystalized ingredient with a clean gauze.

12

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.