Ophthalmic Medicines

 \bullet IRIS 50 is an eye drop that contains neostigmine methylsulfate and vitamin B12 that improve the focus adjustment of the eye, vitamin B2, and taurine that promote metabolism. If has an excellent effect on such symptoms as blurry vision and ocular fatigue. If gives soft sensation of use.

Indication

Eyestrain, prevention of eye troubles (after swimming, or to wash out sweat or dust etc.), ophthalmia by ultraviolet rays etc. (snow blindness etc.), blepharitis (running eye), foreign-body feeling by contact lenses, itchy eyes, blurred vision (eye mucus)

Dosage and administration

Instill 2 to 3 drops per dose 3 to 6 times daily.

Comply with the prescribed dosage and administration instructions.

When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.

Do not touch the tip of the container with the eyelid or eyelash. Do not use the drug when the solution becomes turbid. Do not use the drug when you are wearing soft contact lenses.

Use this only as an eye drop.

ingredient and amount

Neostigimine methylsulfate 0.005% Cyanocobalamin (Vitamin B12) 0.02% Flavin adenine dinucleotide sodium 0.05% Taurine (Aminoethyl Sulfonic Acid) 1% Chondroitin Sulfate Sodium 0.5%

Excipients

I-Menthol, Dryobalanops aromatica resin, Eucalyptus oil, Chlorobutanol, Benzalkonium chloride, citric acid, Sodium citrate, Boric acid, pH adjusting agent, Edetate sodium, Isotonic agents, Polysorbate 80, Anhydrous ethanol

Precautions

Consultation

• The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use. Patients undergoing medical treatment from a physician

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

Severe eye pain

Persons diagnosed as having the following

Glaucoma

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation. Skin

rash/redness

itching

Eyes:

congestion of the eye

itching

swelling

• In case of the following, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

When blurred vision does not improved.

When symptoms do not improve even after using the medicine for about 2 weeks

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with an air tight closure on.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)

Do not share the drug with other people.

Do not leave the drug under high temperatures, such as on a car dashboard. (It may result in the deformation of the container and the degradation of the drug solution.)

If the drug solution stains your clothes while administering it in your eyes, wash with water immediately.

Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).

[Disclaimer on Multilingual OTC Product Information]

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.