

IRIS AG GUARD

Class 2 OTC drug

Eye drops

◆ IRIS AG GUARD contains 3 active ingredients, inhibits the growth of substances that cause allergic symptoms (histamine etc.), alleviates inflammation such as blood shot, and improves the condition of the eye damaged by inflammation. ■ The eye drop gives cool sensation when administered.

Indication

To alleviate the following allergic eye symptoms due to pollen or house dust (indoor dust): blood shot eyes, itching in the eye, blurred vision (due to excessive eye mucus), watery eye, foreign-body sensation (irritating)

Dosage and administration

Instill 1 to 2 drops per dose 4 times daily (morning, noon, evening, and bedtime).

Comply with the prescribed dosage and administration instructions.

When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.

Immediately wipe off the solution that overflowed in areas around the eye.

Be careful; the drug solution may be contaminated or become turbid due to eye mucus and bacteria if the top of the container touches the eyelid or eyelash.

Do not use when the drug appears cloudy.

Do not use the drug as a fitting liquid for contact lenses. Do not use the drug while wearing contact lenses.

Use this only as an eye drop.

For ocular symptoms of seasonal allergy caused by pollen, it is effective to start the drug at the early stage of the symptoms (immediately after minor symptoms start) when pollen begins to scatter.

ingredient and amount

In 100 mL

Ketotifen Fumarate 69mg (50mg as Ketotifen)

Dipotassium Glycyrrhizinate 250mg

Taurine (Aminoethyl Sulfonic Acid) 1000mg

Excipients

I-Menthol,dl-Camphor,Chlorobutanol ,D-Mannitol ,Boric acid,Tromethamine ,Edetate sodium,Benzalkonium chloride

Precautions relating to ingredient and amount

No need to worry if you taste sweetness in the mouth when administering the drug. The sweetness is caused by dipotassium glycyrrhizinate, an active ingredient.

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

● This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Infants under 1 year old.

If the drug is used concurrently with nose drops, do not drive a car or operate machinery after using the drug.

(symptoms such as sleepiness may occur.)

Consultation

● The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.

Patients undergoing medical treatment from a physician

Persons undergoing treatment for allergy, such as desensitization.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

Severe eye pain

Persons who are not sure whether the symptoms are caused either by allergy or other factors.

If you have the following symptoms, which may not be caused by allergies, consult with a physician before use.

Symptoms only in one eye

Ocular symptoms only and no nasal symptoms

Vision is also affected.

Having a lot of eye mucus

Pregnant women or women suspected of being pregnant.

Persons diagnosed as having the following:

Glaucoma

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin
rash/redness
itching
Eyes:
congestion of the eye
Irritation
pain
swelling
itching
blurred vision
dryness of the eyes
Drowsiness
Glare in the eye
headache
Other: dry mouth

*Including the sites around the eyes

**Some people experience symptoms such as severe itching, swelling, and pain in the eyelids, areas around the eyes, and cheeks in the first few days of the use.

● In case of the following, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

When blurred vision does not improve.

Symptoms do not resolve after using the drug for about 1 week.

● In case of the following, consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

The use period exceeds 2 weeks even though the symptoms have been relieved.

Precautions for storage and handling

Avoid direct sunlight, tightly close the cap, and store the drug in a cool place where possible.

Store the product beyond the reach of children.

To prevent misuse and quality deterioration, do not transfer the drug to other containers.

To prevent contamination,

Do not share the drug with other people.

Do not leave the drug under high temperatures, such as on a car dashboard. (It may result in the deformation of the container and the degradation of the drug solution.)

Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).

12

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.