

# ANTIMICROBIAL IRIS

Class 2 OTC drug  
Ophthalmic Medicines

◆ANTIMICROBIAL IRIS is a preservative-free, one-dose eye drop that effectively relieves eye diseases such as sty and conjunctivitis caused by bacteria. No irritation, but soft sensation of use.

## Indication

Conjunctivitis (pink-eye), chalazia, blepharitis (running eye), itchy eyes

## Dosage and administration

Instill 2 to 3 drops per dose 3 to 6 times daily.

Comply with the prescribed dosage and administration instructions.

When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.

Do not touch the tip of the container with the eyelid or eyelash. Do not use the drug when the solution becomes turbid.

Do not use the drug when you are wearing soft contact lenses.

Use this only as an eye drop.

Do not forcibly tear it open. Be careful that the drug solution may be sprinkled if you open the drug forcibly.

This is a one-dose eye drop and does not contain any preservatives. Discard the remaining solution after use.

## ingredient and amount

Sodium Sulfamethoxazole 4%

Dipotassium Glycyrrhizinate 0.1%

## Excipients

pH adjusting agent,Other one-component

## Precautions

### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- Do not take this medicine for a long time

### Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.

Patients undergoing medical treatment from a physician

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

Severe eye pain

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Eyes:

congestion of the eye

itching

swelling

- If symptoms do not improve even after using this medicine for 3 to 4 days, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

## Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)

Do not share the drug with other people.

To prevent the discoloration of the drug solution by light, the drug is contained in a colored container and a colored bag.

Avoid light, and store the drug in a box. Do not store in a refrigerator.

Do not leave the drug under high temperatures, such as on a car dashboard. (It may result in the deformation of the container and the degradation of the drug solution.)

Note that the cap may come off if the drug is placed in the back pocket of pants.

Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.