

TAISHO KAMPO STOMACH MEDICINE <ORAL LIQUID>

Class 2 OTC drug
Gastrointestinal medicine

Indication

Excessive drinking, nausea (churning of stomach due to hangover/drunken sickness, churning of stomach, nausea, vomituration), loss of appetite, dyspepsia, feeling of stomach/abdominal distention, gastrasthenia, overeating, heartburn, heaviness in the stomach, pressure on the chest, and vomiting

Dosage and administration

Adults (15 years or over): Take 1 bottle per dose 3 times daily before a meal or between meals.

Take the drug 2 to 3 hours after a meal when instructed to take the drug between meals.

Comply with the prescribed dosage and administration instructions.

The drug contains herbal extract. Sedimentation occurs over time. This does not affect the drug efficacy. Shake well before taking the drug.

ingredient and amount

In 1 bottle (30 mL)

Goreisanryo Dry Extract 0.3 g 0. 3g (equivalent to 0.833 g of Alisma Tuber,0.750 g of Poria Sclerotium,0.500 g of Cinnamon Bark,0.750 g of Polyporus Sclerotium,0.750 g of Anractylodes Rhizome)

Orengedokuto Dry Extract 0. 3g (equivalent to 0.333g of Coptis rhizome, 0.333g of Phellodendron bark, 0.333g of Gardenia fruit, 0.5g of Scutellaria root)

Excipients

Sugar,D-Sorbitol ,citric acid,Sodium benzoate,Butyl parahydroxybenzoate,Polysorbate 80,Anhydrous ethanol,Other two-component

Precautions

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Mesenteric phlebosclerosis: Abdominal pain, diarrhoea, constipation, abdominal distension, etc. may occur repeatedly with long-term oral administration.

● When symptoms do not improve even after taking the medicine for about 1 month , stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

●If you take this medicine for a long time, consult a physician, pharmacist, or registered salesperson.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not take the product past the expiration date.

The drug contains only one portion of drink. Drink the drug once you open the container. Do not store it after opening the container.

51

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.