

TAISHO KAMPO STOMACH MEDICINE GRANULES

Class 2 OTC drug

◆ Taisho Kampo Ichoyaku is a gastrointestinal medicine containing a combination of Anchusan and Shakuyakukanzoto for people who have weak digestive organs, and have stomach and digestive problems or lose appetite due to irregular eating habits or summer heat. ■ The drug alleviates gastrointestinal symptoms such as heaviness, discomfort of the stomach, and loss of appetite for people who want to enjoy meals, and improve the distressed stomach condition. ■ Particles with good scent and flavor.

Indication

Heavy stomach feeling, feeling of discomfort in the stomach, gastritis, stomach ache, belch, loss of appetite, feeling of abdominal distention, pressure on the chest, heartburn, chlorhydria, abdominal pain, nausea (churning of stomach and vomituration)

Dosage and administration

Take the following dose with cold or lukewarm water before a meal or between meals.

15 years or over: 1 packet per dose, 3 times daily

5 to 14 years: 1/2 packet per dose, 3 times daily

Under 5 years: Do not use

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

ingredient and amount

In 1 packet (1.2 g)

Anchusan 700mg (equivalent to 200mg of Cinnamon bark, 150mg of Corydalis tuber, 150mg of Oyster shell, 75mg of Fennel, 50mg of Amomum seed, 50mg of Glycyrrhiza, 25mg of Ryokyo)

Powdered Shakuyakukanzoto Extract 140mg (Peony root 280mg, Glycyrrhiza 280mg)

Excipients

anhydrous silicic acid, Potato starch, Lactose, cellulose, Hydroxypropylcellulose, Talc

Precautions

Consultation

● The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

● When symptoms do not improve even after taking the preparation for about 1 month, stop taking the medicine and consult a physician (dentist), pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.

Do not take the product past the expiration date.

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.