TAISHO STOMACH MEDICINE K

- ◆TAISHO STOMACH MEDICINE K is a gastrointestinal medicine that contains Shakuyakukanzoto extract and Anchusan.
- ◆TAISHO STOMACH MEDICINE K that contains these 2 herbal medicines relieves gastrointestinal symptoms such as pain and heaviness in the stomach as well as gastritis.

Indication

Stomachache, heavy stomach feeling, gastritis, feeling of discomfort in the stomach, loss of appetite, belch, abdominal pain, heartburn, nausea, chlorhydria, feeling of abdominal distention

Dosage and administration

Take the following dose with cold or lukewarm water before a meal or between meals.

15 years or over: 1 packet per dose, 3 times daily 5 to 14 years: 1/2 packet per dose, 3 times daily

Under 5 years: Do not use

Precaution

- · Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.

Ingredient

In 1 packet (1.2g)

Anchusan 700mg

(equivalent to 200mg of Cinnamon Bark, 150mg of Corydalis Tuber, 150mg of Oyster Shell, 75mg of Fennel, 50mg of Amomum Seed, 50mg of Glycyrrhiza, 25mg of Alpinia officinarum Rhizome)

Shakuyakukanzoto Extract 170mg

(Peony Root 340mg, Glycyrrhiza 340mg)

Excipients

Anhydrous silicic acid, potato starch, lactose, cellulose, hydroxypropylcellulose, talc

Precaution

The drug contains herbal medicine as a raw ingredient. The color of the drug may differ slightly, but this will not affect the efficacy.

Precautions

Consultation

• The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration

Patients undergoing medical treatment from a physician.

The elderly.

Persons with the following symptoms.

Oedema

Persons diagnosed as having the following.

Hypertension, kidney disease, heart disease

• If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Skin: rash/redness, itching

• The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid. Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Pseudoaldosteronism, myopathy: In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress.

22 August 2022 1 / 2page

- When symptoms do not improve after taking this medicine for 1 month, discontinue use of this medicine and consult a physician, pharmacist, or registered salesperson, taking this instruction leaflet with you
- If you take this medicine for a long time, consult a physician, pharmacist, or registered salesperson

Precautions for storage and handling

- · Avoid direct sunlight, and store the product in a cool place with little humidity.
- · Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.
- Do not take the product past the expiration date.

11

【Disclaimer on Multilingual Product Information】

- •This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- •Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- •Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- •The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

22 August 2022 2 / 2page