

STOMATITIS PATCH TAISHO A

Class 3 OTC drug
Stomatitis Products

◆STOMATITIS PATCH TAISHO A is a patch to treat stomatitis and glossitis. It sticks well in the mouth, steadily covers the affected site, while the effect continues for a long time.

Indication

stomatitis, and glossitis

Dosage and administration

Adhere 1 piece per dose onto each mucous membrane of lesion once to 4 times daily.

* Do not use this drug in infants aged under 5 years.

Comply with the prescribed dosage and administration instructions.

The drug does not dissolve. It comes off by itself as time passes. (Do not peel off forcibly.)

This is a patch for the oral mucosa. Do not take it orally.

Use for children \geq 5 years old should be supervised by a parent.

For the use of the drug in children, be careful that they may remove the applied drug with fingers.

The drug may not be properly pasted in wrong usage. Read the instruction for use carefully, and use it properly.

Discard the drug, once coming off, instead of swallowing.

ingredient and amount

In 1 piece (1 patch)

Lithospermum Root Extract 0.10mg (equivalent to 4.55 mg of the crude drug)

Glycyrrhetic Acid 0.15mg

Excipients

Polyacrylic acid, Triethyl citrate, Hypromellose, Ethylcellulose, Castor oil, Titanium oxide

Precautions

Consultation

● The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before use.

Patients undergoing medical treatment from a physician or dentist.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with chronic symptoms

Those with extensive lesions.

● If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, dentist, pharmacist, or registered sales person for a consultation.

Skin

rash/redness

itching

● When symptoms do not improve even after using the medicine 5 to 6 days, stop using this medicine and consult a physician, dentist, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers.

(It may lead to misuse or quality deterioration)

Put the unused drug in the bag, and fold up the opening for storage.

Do not use the product past the expiration date.

Even within the expiration date, use up as soon as possible once it is opened (to assure the quality).

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.