

LIPOVITAN D SUPER

Newly designated quasi-
drugs

Health drinks

Indication

Nutritional supplements during physical fatigue, during and after illness (or when physical strength is impaired after illness), nutritional disorders, feverish wearing disease□

Loss of appetite

For nutritional supplementation during pregnancy/nursing

Nutrients/tonics

Weak constitution

Dosage and administration

Adults (15 years or over): Take 1 bottle (100 mL) per dose once daily.

Under 15 years: Do not take

Comply with the prescribed dosage and administration instructions.

(Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug.)

The drug contains a herbal extract. Sedimentation may occur in rare cases. This does not affect drug efficacy.

Shake well before taking the drug.

ingredient and amount

In 100 mL

Thiamine nitrate 10mg

Riboflavin sodium phosphate 5mg

Pyridoxine hydrochloride (vitamin B6) 5mg

Tocopherol acetate 10mg

Nicotinamide 20mg

Taurine (Aminoethyl Sulfonic Acid) 2000mg

L-Arginine Hydrochloride 300mg

Anhydrous caffeine 50mg

Inositol 100mg

Ginseng extract 130mg

Excipients

Sugar,D-Sorbitol ,Povidone ,Glycerol fatty acid ester,Glycerin,Propyl gallate,Sodium benzoate,pH adjusting agent,citric acid,flavor,Vanillin,,

Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

Precautions

Consultation

In the following cases, immediately discontinue this drug, and consult your physician or pharmacist. Be sure to show this product.

If the following symptoms occur after taking this drug

Skin

rash

Gastrointestinal system

feeling of discomfort in the stomach

When symptoms do not improve even after taking the medicine for a while

The following symptoms may occur. If these symptoms persist or worsen, discontinue this drug, and consult your physician or pharmacist.

diarrhea

Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.

Store the product beyond the reach of children.

Do not take the product past the expiration date.

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.