

LIPOVITAN F i n e PRECIOUS

Newly designated quasi-
drugs

Health drinks

Indication

Nutritional supplements during physical fatigue, during and after illness (or when physical strength is impaired after illness), nutritional disorders, feverish wearing disease□

Loss of appetite

For nutritional supplementation during pregnancy/nursing

Nutrients/tonics

Weak constitution

Dosage and administration

Adults (15 years or over): Take 1 bottle (100 mL) per dose once daily.

Under 15 years: Do not take

Comply with the prescribed dosage and administration instructions.

(Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug.)

The drug contains a herbal extract. Sedimentation may occur in rare cases. This does not affect drug efficacy.

Shake well before taking the drug.

ingredient and amount

In 100 mL

Taurine (aminoethyl sulfonic acid) 1500mg

Royal jelly 100mg

Whitethorn extract S 12.5mg

Lycium Fruit Fluid Extract 0.2mL

Thiamine nitrate 10mg

Riboflavin sodium phosphate 5mg

Pyridoxine hydrochloride (vitamin B6) 5mg

Nicotinamide 20mg

Inositol 100mg

Anhydrous caffeine 50mg

Excipients

Erythritol ,Acesulfame K,Sucralose ,Stevia extract,citric acid,Sodium citrate,Povidone ,Propyl gallate,Sodium benzoate,flavor,Vitamin E ,D-Sorbitol ,Glycerin,l-Menthol,Alcohol (not more than 0.15 mL),,

Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

Precautions

Consultation

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash

Gastrointestinal system

feeling of discomfort in the stomach

The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

diarrhea

When symptoms do not improve even after taking the medicine for a while, stop taking this medicine, and show this product to a physician, pharmacist, or registered salesperson for consultation.

Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.

Store the product beyond the reach of children.

Do not take the product past the expiration date.

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.