

LIPOVITAN DX PLUS

- ◆ LIPOVITAN DX PLUS contains vitamin B₁₂ and three herbal medicines in addition to taurine and vitamins B₁, B₂ and B₆, which are the base components of the Lipovitan series, which has brought relief to fatigue in the lifestyles of many. With just one dose a day, this medicine is highly effective against the "fatigue" felt in daily life.
- ◆ It has an effect that improves and prevents tiredness of the eyes due to poor nutrition.
- ◆ It has an effect that improves and prevents problems of the shoulders, neck, low back or knees associated with aging.
- ◆ This product is recommended for middle-aged and elderly people who feel easily fatigued and problems of the shoulders, neck and low back associated with aging, in order to stay vigorous and active every day.

Indication

- ★ Recovery from and prophylaxis of fatigue
- ★ Improvement and prophylaxis of physical malfunction associated malnutrition in daily living
 - Eyestrain
 - Bad conditions of the shoulders, neck, lower back or knees
 - Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid
 - Being sensitive to the cold, poor blood circulation
 - Bad skin conditions (rough or dry skin)
 - Decreased appetite and malaise associated with hangover
- ★ Improvement and prophylaxis of physical malfunction associated with frail constitution (including physical weakening due to aging.)
 - Bad conditions of the shoulders, neck, lower back or knees
 - Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid
 - Being sensitive to the cold, poor blood circulation
 - Bad skin conditions (rough or dry skin)
- ★ Maintenance and improvement of physical strength or resistance or concentration
- ★ Nutritional support for declined physical strength during/after illness, febrile debilitating illness, anorexia, pregnancy/lactation period, before and after childbirth, etc.

Dosage and administration

Take the following dose with cold or lukewarm water.

Adults (15 years or over): 3 tablets per dose, 1 times daily

Under 15 years: Do not use

Precaution

- Comply with the prescribed dosage and administration instructions. (Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug)
- Do not return the tablets touched by wet fingers into the jar. (This causes discoloration and may deteriorate tablet quality)
- The small package wrapped with paper in the jar is a drying agent. Do not take it.

Ingredient

In 3 tablets

Taurine (aminoethyl sulfonic acid) 500mg

Thiamine nitrate (vitamin B₁) 10mg

Riboflavin (vitamin B₂) 5mg

Pyridoxine hydrochloride (vitamin B₆) 5mg

Cyanocobalamin (vitamin B₁₂) 60 μg

Lycium Fruit Fluid Extract-A 0.2mL (equivalent to 200mg of the crude drug)

Fluid extract of Eucommia leaf 60 μL (equivalent to 60mg of Eucommia leaves)

Japanese Angelica Root extract S 30mg (equivalent to 120mg of Japanese Angelica Root)

Excipients

Anhydrous silicic acid, cellulose, hydroxypropylcellulose, magnesium stearate, talc, macrogol, titanium oxide, hypromellose, ferric oxide, carnauba wax

Precaution

- The drug contains herbal medicine as a raw ingredient. The color of the drug may differ slightly. This does not impact its efficacy.
- No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B₂ contained in the drug.

Precautions

Consultation

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
 - Skin: rash
 - Gastrointestinal system: feeling of discomfort in the stomach
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
 - Diarrhea
- When symptoms do not improve even after taking this medicine for a while, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you

Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight in an air-tight container.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened. (to assure the quality)

11

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.