

LIPOVITAN ACTIVE

Designated Quasi-drug

Precautions for Use

Things To Consult On

1. **If the following symptoms appear after drinking, you may be experiencing side effects. Discontinue use immediately and consult with your doctor, pharmacist or registered seller with this product to hand.**
Skin: rash / Digestion: stomach discomfort
2. **The following symptom may appear after drinking. If this symptom persists or worsens, discontinue use and consult with your doctor, pharmacist or registered seller with this product to hand.**
Diarrhea
3. **If symptoms do not improve after a while, discontinue use and consult with your doctor, pharmacist, or registered seller with this product to hand.**

Efficacy

★ Relieves and prevents fatigue ★ Improves and prevents physical problems associated with a weakened constitution (including physical infirmities associated with aging): tiredness, persistent tiredness, lack of physical stamina, lethargy, sluggishness, sensitivity to cold, poor blood circulation, shoulder/neck/lower back/knee discomfort, poor complexion (rough or dry skin), gastrointestinal weakness causing frequent diarrhea, bone or dental deterioration ★ Improves and prevents physical problems associated with malnourishment in daily life: decreased appetite and sluggishness associated with hangover, eye fatigue ★ Maintains and improves physical stamina, resilience, and concentration ★ Supplements nutrition when stamina is reduced during and after illness, when suffering from a debilitating illness with fever, when experiencing loss of appetite, or during lactation or before and after childbirth

Usage/Dosage

Adults (15 years or over): Drink one bottle (50 mL) once daily.

*Under 15 years: Do not drink.

Note:

- (1) Strictly comply with the prescribed usage/dosage instructions. (Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this product.)
- (2) As this product contains herbal extract, sedimentation may occur in rare cases. This does not affect drug efficacy. Shake well before drinking.

Ingredients in one medium bottle (50 mL)

Taurine (2-aminoethyl sulfonic acid).....	1500 mg	Thiamine nitrate (vitamin B ₁).....	5 mg
Royal jelly.....	200 mg	Riboflavin sodium phosphate (vitamin B ₂) ...	3 mg
Ginger fluid extract	0.3 mL	Pyridoxine hydrochloride (vitamin B ₆)	5 mg
(Equivalent to 300 mg of ginger)		Nicotinamide	20 mg
Sun-dried tangerine orange peel extract	20 mg	Inositol.....	50 mg
(Equivalent to 100 mg of dried peel)		Caffeine anhydrous.....	50 mg
Angelica root fluid extract S.....	0.2 mL		
(Equivalent to 200 mg of angelica root)			

Additives: White sugar, D-sorbitol, citric acid, sodium citrate, povidone, sodium benzoate, flavoring, alcohol, glycerin (not more than 0.20 mL alcohol)

Note:

Do not worry if your urine is yellow after drinking the product. This is caused by vitamin B₂ contained in the product.

Precautions for Storage and Handling

- (1) Store the product in a cool place that is out of direct sunlight.
- (2) Please store this product outside of the reach of children.
- (3) Do not take the product after the expiration date.

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.