

LIPOVITAN GOLD ACE

Class 2 OTC drug

Precautions for Use

Things To Consult On

1. If the following symptoms appear after drinking, you may be experiencing side effects. Discontinue use immediately and consult with your doctor, pharmacist or registered seller with this product to hand.
Skin: Rash/Digestive system: Stomach discomfort
2. The following symptom may appear after drinking. If this symptom persists or worsens, discontinue use and consult with your doctor, pharmacist or registered seller with this product to hand.
Diarrhea
3. If symptoms do not improve after a while, discontinue use and consult with your doctor, pharmacist, or registered seller with this product to hand.

Efficacy:

★ Nourishing and strengthening ★ Weak constitution ★ Physical fatigue, illness recovery, loss of appetite, nutritional disorders, febrile debilitating illnesses, pregnancy and lactation, etc.

Usage/Dosage:

Adults (15 years or over): Drink 1 bottle (50 mL) once daily.

Note: Do not take if under age 15.

Note:

- (1) Strictly follow the specified dosage and administration.
- (2) As this product contains herbal extract, sedimentation may occur in rare cases. This will not affect the efficacy of the product. Shake well before drinking.

Ingredients in one medium bottle (50 mL)

Taurine (2-aminoethyl sulfonic acid)	1500 mg	Riboflavin sodium phosphate (vitamin B ₂) ...	5 mg
Royal jelly	200 mg	Pyridoxine hydrochloride (vitamin B ₆)	5 mg
Carrot extract-B.....	90 mg	Tocopherol acetate (vitamin E)	30 mg
(Equivalent to 630 mg of carrot)		Nicotinamide	20 mg
Astragalus root extract-C.....	0.2mL	Inositol.....	100 mg
(Equivalent to 200 mg of astragalus)		Caffeine anhydrous	50 mg
Molindae root extract.....	0.2mL		
Equivalent (Equivalent to 200 mg of morinda officinalis)			
Thiamine nitrate (vitamin B ₁).....	5 mg		
Additives: Sugar, D-sorbitol, sucralose, acesulfame-K, povidone, glycerin fatty acid ester, polyoxyethylene hydrogenated castor oil, glycerin, sodium benzoate, citric acid, sodium citrate, flavoring, alcohol, vanillin, vitamin E (0.2 mL or less of alcohol)			

Note:

Do not worry if your urine is yellow after taking this product. This is caused by the vitamin B₂ contained in the product.

Precautions for Storage and Handling:

- (1) Store the product in a cool place that is out of direct sunlight.
- (2) Please store this product outside of the reach of children.
- (3) Do not take the product after the expiration date.

45 kcal per bottle

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.