

LIPOVITAN Jr.

Class 3 OTC drug
Vitamin-containing health
drugs

Indication

☆ Nutritional support for growth in children, unbalanced diet in children, febrile debilitating illness, during/after illness, loss of appetite, malnutrition, etc.

Nutrients/tonics

Weak constitution

Dosage and administration

8 to 14 years: Take 1 bottle (50 mL) per dose once daily.

Under 8 years: Do not take

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

The drug contains a herbal extract. Sedimentation may occur in rare cases. This does not affect drug efficacy.

Shake well before taking the drug.

ingredient and amount

In 50 mL

Schisandra Fruit Fluid Extract 0.2mL (equivalent to 200 mg of the crude drug)

Eleutherococcus Senticosus Rhizoma Dry Extract 16mg (equivalent to 400 mg of the crude drug)

Calcium Gluconate Hydrate 123.75mg

Calcium chloride 55.5mg

Taurine (Aminoethyl Sulfonic Acid) 800mg

Carnitine Hydrochloride 50mg

Magnesium L-aspartate 100mg

Thiamine Nitrate 4mg

Riboflavin sodium phosphate 4mg

Pyridoxine Hydrochloride (vitamin B6) 4mg

Nicotinamide 15mg

Excipients

Sugar,Reduced maltose syrup,Povidone ,Benzoic acid,pH adjusting agent,flavor,Glycerin

Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

Precautions

When not to use the product

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash

Gastrointestinal system

feeling of discomfort in the stomach

The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

diarrhea

When symptoms do not improve even after taking the medicine for a while, stop taking this medicine, and show this product to a physician, pharmacist, or registered salesperson for consultation.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not take the product past the expiration date.

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.