

Read the instruction sheet in this box carefully before using.

LIPOVITAN DX M

Designated Quasi-drug



Precautions for Use:

Things To Consult On




1. If the following symptoms appear after drinking, you may be experiencing side effects. Discontinue use immediately and consult with your doctor, pharmacist or registered seller with this box to hand.
Skin: Rash/Digestive system: Stomach discomfort
2. The following symptom may appear after drinking. If this symptom persists or worsens, discontinue use and consult with your doctor, pharmacist or registered seller with this box to hand.
Diarrhea
3. If symptoms do not improve after a while, discontinue use and consult with your doctor, pharmacist, or registered seller with this box to hand.

Efficacy:

★ Relieves and prevents fatigue ★ Improves and prevents physical ailments associated with a weak constitution (including physical weakness due to aging): muscle weakness, fatigability, lingering fatigue, lack of stamina, feeling heavy or sluggish, shoulder/neck/back/knee discomfort, bone or tooth weakness
★ Maintenance and improvement of stamina, physical endurance or concentration ★ Improvement and prevention of physical discomfort due to poor nutrition in everyday life: muscle weakness, fatigability, lingering fatigue, lack of stamina, feeling heavy or sluggish, shoulder/neck/back/knee discomfort, ★ Supplemental nutrition for when physical stamina is reduced during and after illness, when suffering from a debilitating illness accompanied by fever, when experiencing a loss of appetite, or during lactation or before and after childbirth

Usage/Dosage:

Take the following dosages with room-temperature or lukewarm water.

Age	Single dose	Number of doses
Adults (15 years and older)	3 tablets 	Once per day
Under 15 years old	Do not take this medicine	

Note:

- (1) Strictly comply with the prescribed usage/dosage instructions. (Pay attention to excessive intake, etc., if a product containing other vitamins, etc., is taken together with this product.)
- (2) Do not put tablets that have been touched with wet hands back in the bottle. (This may result discoloration or a change in quality)
- (3) Do not ingest the desiccant inside the bottle, which is wrapped in wrapping paper.

Ingredients: In 3 tablets

Taurine (2-aminoethyl sulfonic acid).....	200 mg	L-isoleucine.....	50 mg
Thiamine nitrate (vitamin B ₁)	10 mg	L-arginine hydrochloride	30 mg
Riboflavin (vitamin B ₂)	5 mg	Magnesium L-aspartate.....	180 mg
Pyridoxine hydrochloride (vitamin B ₆).....	5 mg	L-histidine hydrochloride hydrate	6 mg
L-valine.....	50 mg	Anhydrous calcium hydrogen phosphate ...	102 mg
L-leucine.....	100 mg	(30 mg as calcium)	

Additives: silica, cellulose, hydroxypropyl cellulose, magnesium stearate, talc, macrogol, titanium oxide, hypromellose, ferric oxide, carnauba wax

Note:

Do not worry if your urine is yellow after drinking the product. This is caused by vitamin B₂ contained in the product.

Precautions for Storage and Handling:

- (1) Store sealed in a cool, dry place away from direct sunlight.
- (2) Please store this product outside of the reach of children.
- (3) Do not transfer to another container. (This may result in misuse or a change in quality)
- (4) Do not take the product after the expiration date. Use this medicine within 6 months of opening, even if still within the expiration date. (In order to ensure quality.)

<The first time you take this medicine>

- ① Peel off the bottle cap seal and discard it.
- ② Remove and discard the filling. (This filling is intended to prevent damage to capsules during transport. Removing and inserting this repeatedly may cause foreign matter contamination)

– This text is the information on the packaging (product box) –

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.