

# LIPOVITAN CLEAR

Newly designated quasi-  
drugs

Health drinks

## Indication

- ☆ Maintenance and improvement of physical strength or resistance or concentration
- ☆ Recovery from and prophylaxis of fatigue
- ☆ Improvement and prophylaxis of physical malfunction associated with frail constitution (including physical weakening due to aging): weakening of bones or teeth
- ☆ Improvement and prophylaxis of physical malfunction associated malnutrition in daily living:  
Bad conditions of the shoulders, neck, lower back or knees  
Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid  
Bad skin conditions (rough or dry skin)  
Decreased appetite and malaise associated with hangover
- ☆ Nutritional support for declined physical strength during/after illness, febrile debilitating illness, anorexia, pregnancy/lactation period, before and after childbirth, etc.

## Dosage and administration

Adults (15 years or over): Take 1 bottle (100 mL) per dose once daily.  
Under 15 years: Do not use  
Comply with the prescribed dosage and administration instructions.  
(Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug.)

## ingredient and amount

In 1 bottle (50 mL)  
Taurine (aminoethyl sulfonic acid)  
1500mg  
Lycium Fruit Fluid Extract-A  
0.2mL  
(equivalent to 200 mg of the crude drug)  
Royal jelly  
300mg  
Carnitine hydrochloride  
50mg  
Acanthopanax Fluid Extract 0.2mL  
(equivalent to 200 mg of the crude drug)  
Thiamine nitrate (vitamin B1)  
10mg  
Riboflavin  
5mg  
Pyridoxine hydrochloride  
10mg  
Nicotinamide  
30mg  
Inositol  
100mg  
Anhydrous caffeine  
50mg

## Excipients

Sugar, D-Sorbitol, Sucralose, Povidone, Sodium benzoate, Sodium citrate, Citric acid, pH adjusting agent, Flavoring, alcohol, Glycerin, Vanillin, (Alcohol not more than 0.2 mL)

## Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

## Precautions

### Consultation

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

Skin  
rash  
Gastrointestinal system

feeling of discomfort in the stomach

The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

diarrhea

When symptoms do not improve even after taking the medicine for a while, stop taking this medicine, and show this product to a physician, pharmacist, or registered salesperson for consultation.

### Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.

Store the product beyond the reach of children.

Do not take the product past the expiration date.

11

#### 【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.