

LIPOVITAN CLEAR

Designated Quasi-drug

Precautions for Use

Things To Consult On

1. **If the following symptoms appear after drinking, you may be experiencing side effects. Discontinue use immediately and consult with your doctor, pharmacist or registered seller with this product to hand.**
Skin: rash / Digestion: stomach discomfort
2. **The following symptom may appear after drinking. If this symptom persists or worsens, discontinue use and consult with your doctor, pharmacist or registered seller with this product to hand.**
Diarrhea
3. **If symptoms do not improve after a while, discontinue use and consult with your doctor, pharmacist, or registered seller with this product to hand.**

Efficacy

☆ Maintenance and improvement of physical strength, resistance or concentration ☆ Recovery from and prophylaxis of fatigue ☆ Improvement and prophylaxis of physical malfunction associated with frail constitution (including physical weakening due to aging): Weakening of bones or teeth ☆ Improvement and prophylaxis of physical malfunction associated malnutrition in daily living: Bad conditions of the shoulders, neck, lower back or knees; Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid; Bad skin conditions (rough or dry skin); Decreased appetite and malaise associated with hangover; Eyestrain ☆ Nutritional support for declined physical strength during/after illness, febrile debilitating illness, anorexia, pregnancy/lactation period, before and after childbirth, etc.

Usage/Dosage

Adults (15 years or over): Drink one bottle (50 mL) once daily.

*Under 15 years: Do not drink.

Note:

- (1) Strictly comply with the prescribed usage/dosage instructions. (Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this product.)
- (2) As this product contains herbal extract, sedimentation may occur in rare cases. This does not affect drug efficacy. Shake well before drinking.

Ingredients in one medium bottle (50 mL)

Taurine (2-aminoethyl sulfonic acid).....	1500 mg	Thiamine nitrate (vitamin B ₁).....	10 mg
Lycium fruit fluid extract-A: 0.2 mL		Sodium riboflavine phosphate (vitamin B ₂)..	5 mg
(equivalent to 200 mg lycium)		Pyridoxine hydrochloride (vitamin B ₆)	10 mg
Royal jelly	300 mg	Nicotinamide	30 mg
Carnitine chloride	50 mg	Inositol.....	100 mg
Acanthopanax fluid extract	0.2 mL	Caffeine anhydrous.....	50 mg
(equivalent to 200 mg of acanthopanax)			
Additives: Sugar, D-sorbitol, sucralose, povidone, sodium benzoate, sodium citrate, citric acid, pH adjusting agent, flavoring, alcohol, glycerin, vanillin			
(alcohol not more than 0.2 mL)			

Note:

Do not worry if your urine is yellow after drinking the product. This is caused by vitamin B₂ contained in the product.

Precautions for Storage and Handling

- (1) Store the product in a cool place that is out of direct sunlight.
- (2) Please store this product outside of the reach of children.
- (3) Do not take the product after the expiration date.

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.