

# HONZO YOKUININ TABLETS S

## Indication

Warts, rough skin

## Dosage and administration

Take the following dose with cold or lukewarm water before a meal or between meals.

15 years or over: 6 tablets per dose, 3 times daily

11 to 14 years: 4 tablets per dose, 3 times daily

8 to 10 years: 3 tablets per dose, 3 times daily

5 to 7 years: 2 tablets per dose, 3 times daily

Under 5 years: Do not take

## Precaution

- Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.
- Do not return the tablets touched by wet fingers into the jar. (It will cause discoloration and may deteriorate the quality.)

## Ingredient

In a daily dose (18 tablets)

Powdered coix seed 4,200mg

## Excipients

Silicon dioxide

## Precaution

The drug contains herbal medicine as a raw ingredient. The color of the drug may differ slightly, but this will not affect the efficacy.

## Precautions

### Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
  - Patients undergoing medical treatment from a physician
  - Pregnant women or women suspected of being pregnant.
  - Patients who have experienced allergic symptoms associated with drugs, etc.
- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.
  - Skin: rash/redness, itching
  - Gastrointestinal system :feeling of discomfort in the stomach
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen , discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.
  - diarrhea
- When symptoms do not improve even after taking the preparation for about 1 month, discontinue use and consult a physician, pharmacist or registered salesperson, taking this document with you.

## Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)
- Do not take the product past the expiration date. Even within the expiration date, try to take it up as soon as possible once it is opened (to assure the quality).

**【Disclaimer on Multilingual Product Information】**

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.