

HONZO DAISAIKOTO EXTRACT TABLETS H

Indication

For those who have full physical strength, but suffer from pain in the area from the flank over to the upper abdomen, and tend to develop constipation and have the following symptoms: Gastritis; habitual constipation; stiff shoulders, headache, and constipation associated with hypertension or obesity; neurosis; and obesity disease

Dosage and administration

Take the following dose with cold or lukewarm water before a meal or between meals.

15 years or over: 4 tablets per dose, 3 times daily

5 to 14 years: 3 tablets per dose, 3 times daily

Under 5 years: Do not use

Take the drug 2 to 3 hours after a meal when instructed to take the drug between meals.

Precaution

- Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.
- Do not return the tablets touched by wet fingers into the jar. (It will cause discoloration and may deteriorate the quality)
- Take out and discard the stuffing when you take the first dose.
(This stuffing is to prevent tablets from damage during transport. If the stuffing is taken in and out of the bottle, it may cause contamination)

Ingredient

In 12 tablets

Daisaikoto extract (1/2 portion) 1.70g

Extracted from (Bupleurum Root 3.00g, Rhizoma Pinelliae 2.00g, Scutellaria Root 1.50g, Peony Root 1.50g, Jujube 1.50g, Immature Orange 1.00g, Rhubarb 1.00g, Ginger 0.75g)

Excipients

Cellulose, calcium carbonate, magnesium aluminometasilicate, croscarmellose sodium, magnesium stearate

Precaution

The drug contains herbal medicine as a raw ingredient. The color of the drug may differ slightly, but this will not affect the efficacy.

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur)

- This medicine should not be taken together with the following medicines
Other cathartic agents
- For those who are lactating, do not use this medicine or avoid lactating while using this medicine

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration
Patients undergoing medical treatment from a physician.
Pregnant women or women suspected of being pregnant.
Persons who is physically weak (declining of physical strength or constitutional weakness).
Persons with weak stomach and intestine who have a tendency to diarrhea, diarrhea with severe abdominal pain.
Patients who have experienced rash/redness, itching, etc. with drugs, etc.
- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
Skin: rash/redness, and itching
Gastrointestinal system: diarrhea with severe abdominal pain, abdominal pain

- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.
Interstitial pneumonia: shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance.
Hepatic function failure: symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
Diarrhea
- When symptoms persist after taking this medicine for about 1 month (5-6 days for habitual constipation, or constipation associated with hypertension or obesity), discontinue use, and consult with a physician, pharmacist, or registered salesperson with this explanatory document

Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight in an air-tight container.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not take the product past the expiration date. Even within the expiration date, try to finish it as soon as possible once it is opened. (to assure quality)

11

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.