

◆Meditreat is a drug to treat recurrent vaginal candidiasis with miconazole nitrate as main ingredient. ◼Miconazole nitrate treats vaginal candidiasis by killing Candida, the causative agent of vaginal candidiasis. ◼ The drug relieves vaginal discharge/itching in the sensitive area caused by recurrent vaginal candidiasis.

## Indication

Recurrent vaginal candidiasis (limited to persons who have previous diagnosis/treatment by a physician)

## Dosage and administration

Aged  $\geq 15$  years old and  $< 60$  years old: 1 dose: once daily (before bed time)

Children under 15 years old and persons  $\geq 60$  years old: Do not use the product.

For an adult (aged  $\geq 15$  years old and  $< 60$  years old), insert a suppository deep inside the vagina once daily before going to bed. Use the suppository continuously for 6 days.

If symptoms are not relieved after using the drug for 3 days or if symptoms are not resolved after using the drug for 6 days, seek treatment from a physician.

Take a bath or wash your vaginal area with warm water before use.

Comply with the prescribed dosage and administration instructions.

If the drug is too soft, cool it for a while before use.

If the drug is too hard, leave the drug in room temperature for a while, and use it after it gets soft enough.

Use only in the vagina.

Wash your hands well before and after use.

Do not use an applicator. (Not good for hygiene.)

Even if symptoms are resolved, use the drug for 6 days from the start of use.

Do not use the drug during your period. If your period starts during the use, stop using the drug. In such a case, seek treatment from a physician to confirm whether the symptoms have resolved. (During a period, the drug may leak, and sufficient effects may not be achieved.)

## ingredient and amount

In 1 piece

Miconazole nitrate 100mg

## Excipients

Hard fat

## Precautions

### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

This product should not be used in the following persons:

Persons with seemingly first onset (Other diseases may be the cause in persons with first onset and may require diagnosis by a physician.)

Persons who have had an allergic symptom to the drug or the drug ingredients. (Allergic symptoms may recur in the use of this drug.)

Children under 15 years old or persons  $\geq 60$  years old. (The symptom in children under 15 years old is highly likely the first onset. The symptom in persons  $\geq 60$  years old may be a mixed infection caused by other disease or other bacteria.)

Pregnant women or women suspected of being pregnant. (Take thorough precaution in using the drugs, and consult with a specialist doctor for instructions.)

Persons with fever, chill, lower abdominal pain, shoulder and back pain, colored or bloody vaginal discharge, fishy vaginal discharge, menstruation stopped, irregular or abnormal vaginal bleeding, ulcer, edema, or sores in the vagina or vulva (This may possibly be other diseases that require diagnosis by a physician.)

Persons diagnosed as having the following:

Diabetes (The disease may be frequently repeated, thus requiring diagnosis by a physician.)

Persons on anticoagulants, such as warfarin. (Bleeding tendency, which is an action of warfarin, may be enhanced.)

Persons repeating this disease frequently (once in 1 or 2 months or at least twice in 6 months).

Recurrent vaginal candidiasis is suspected. (If it is difficult to assess by yourself, seek diagnosis from a physician.)

Do not use in the following sites.

Sites other than the vagina (The drug is effective only for Candida infection in the vagina.)

Do not use the following drugs for the vulva when using this drug.

Dermatologic agents other than drugs for candida (These agents may worsen the symptoms or delay the cure.)

## Consultation

The following persons should contact a physician or pharmacist for a consultation before using the drug:

Patients undergoing medical treatment from a physician

(The drug may affect the action of other drugs prescribed by a physician.)

Persons who have had an allergic symptom to drugs.

(Persons who had allergic reaction to a drug may be allergic to the drug.)

Nursing women (Thorough precaution is required for the use of the drug.)

The following symptoms may occur after using the drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to a physician or pharmacist for a consultation.

Other than vagina: urticaria and itching

The following symptoms may occur after use. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to a physician or pharmacist for a consultation.

Vagina: itching, redness, pain, feeling hot, and irritation

(They are considered allergic symptoms to the drug or powerful pharmacological actions of the drug. In such cases, the continuous use of the drug may worsen the symptoms.)

If the symptoms do not improve after using the drug for 3 days or if the symptoms do not resolve after using the drug for 6 days, seek medical treatment from a physician. (The symptoms may be severe or caused by other diseases.)

## Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity and temperature below 30°C. (Given that the drug dissolves at body temperature, store it in a refrigerator after purchasing.)

Store the product beyond the reach of children.

For storage, store in the outer box by placing the tip downward and stand them according to the instruction on the outer box (to prevent the deformation of the suppositories).

Do not touch the product with contraceptive latex products, such as condoms and pessaries. (The contraceptive products may be deteriorated or damaged.)

Do not use the product past the expiration date (to assure quality).

32

### 【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.