

NARON GRANULES

Designated Class 2 OTC
drug

Antipyretic analgesics

◆NARON GRANULES is an antipyretic analgesic to relieve headache, toothache and menstrual pain. Antipyretic analgesic effect of acetaminophen and ethenzamide, and sedative effect of bromovalerylurea effectively relieve these pains. ■This is an antipyretic analgesic that does not contain pyrine medicine.

Indication

1) Relief of headache, toothache, pain after tooth extraction, sore throat (throat pain), earache, joint pain, neuralgia, lumbago, muscular pain, pain due to stiff shoulders, contusion pain, bone fracture pain, pain associated with sprain (sprain pain), painful menses (menstrual pain), and traumatic pain
2) Relief of fever at the time of chills (feeling cold due to fever) and fever

Dosage and administration

Take the following dose with cold or lukewarm water. Avoid taking the drug on an empty stomach whenever possible. Provide a dosing interval for at least 4 hours.

15 years or over: 1 packet per dose, up to 3 times daily

11 to 14 years: 2/3 packet per dose, up to 3 times daily

7 to 10 years: 1/2 packet per dose, up to 3 times daily

3 to 6 years: 1/3 packet per dose, up to 3 times daily

1 to 2 years: 1/4 packet per dose, up to 3 times daily

Under 1 year: Do not take

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

ingredient and amount

In 1 packet (1.6 g)

Acetaminophen 265mg

Ethenzamide 300mg

Bromvalerylurea 200mg

Anhydrous caffeine 50mg

Excipients

Lactose ,Carboxymethylcellulose calcium ,Hydroxypropylcellulose,Potato starch,Magnesium aluminometasilicate ,l-Menthol

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Persons who experienced asthma after taking this drug, other antipyretic analgesics, or cold medicine.

● This medicine should not be taken together with any of the following medicines:

Other antipyretics/analgesics, cold medicine, sedatives, medicine for motion sickness

●After taking this drug, do not drive a car or operate machinery (sleepiness may occur).

Do not drink alcohol before and after taking this drug.

● Do not take this medicine for a long time

Consultation

●The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Baby/infants/children (under 15 years old) who have varicella (chickenpox) or influenza or are suspected of having either of them.

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons diagnosed as having the following

heart disease

kidney disease

Liver disease

Gastric/duodenal ulcer

●If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a

consultation.

Skin

rash/redness

itching

Gastrointestinal system

Vomiting

Vomiting

Loss of appetite

Neuropsychiatric system

dizziness

Other:

Excessive temperature decrease

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joint...)

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.

Do not take the product past the expiration date.

61

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.