•Revised April 2026
(Revised the Precautions for Use)

Make sure to read this instruction sheet before taking this medicine. Additionally, store this sheet so that it can be referenced when needed.

Contains
Acetaminophen
Ethenzamide

For headache and fever

NARON GRANULES

Class 2 OTC drug

<Antipyretic analgesic>

◆ NARON GRANULES are an antipyretic analgesic for those suffering from headache, toothache, menstrual pain, or other such pain. They help to effectively relieve these pains through the antipyretic and analgesic effects of acetaminophen and ethenzamide, and the sedative effects of bromovalerylurea.
◆ This is an antipyretic analgesic drug that does not contain pyrazines.

Precautions for Use

Things You Must Not Do

(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects or accidents occurring)

1 The following persons should not take this medicine:
(1) Persons who have had allergic symptoms as a result of taking this medicine or any component of it.
(2) Persons who have experienced asthma while taking this medicine or other antipyretic analgesics or cold medicines.

2 Do not take any of the following medicines while taking this medicine:
Other antipyretic analgesics, cold medicines, sedatives, medicines for motion sickness

3 Do not drive a vehicle or operate machinery after taking this medicine
(You may experience drowsiness, etc.)

4 Do not drink alcohol before or after taking this medicine

5 Do not use for long periods of time

6 Do not take this medicine in excess amount.*

Things To Consult On

1 The following individuals should consult with their doctor, dentist, pharmacist or a registered seller before taking this medicine:
(1) Persons receiving treatment from a doctor or dentist.
(2) Persons who are pregnant or who think they may be pregnant.
(3) Infants, toddlers and children (under 15) who have or are suspected of having chickenpox or influenza.
(4) Elderly persons.
(5) Persons with a history of allergic reactions caused by medicines, etc.
(6) Persons who have received any of the following diagnoses:
Heart disease, kidney disease, liver disease, stomach or duodenal ulcer

2 If any of the following symptoms appear after you have taken this medicine, you may be experiencing side effects. As such, discontinue use immediately and consult with your doctor, pharmacist or registered seller, bringing this instruction sheet with you

Relevant Area	Symptoms
Skin	Rash, redness, itching
Digestive system	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Other	Excessive hypothermia

In rare cases, the serious side effects listed on the right may occur. In such cases, see a doctor immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Itching, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion immediately after taking this medicine.
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	High fever, bloodshot eyes, eye discharge, lip sores, sore throat, widespread skin rash/redness, small bumps (small pustules) appearing on reddened skin, whole body sluggishness, loss of appetite, etc., that persists or worsens rapidly.
Drug-induced hypersensitivity syndrome	Widespread redness of the skin, generalized rash, fever, fatigue, swelling of the lymph nodes (e.g. neck, armpits, groin), etc.
Liver dysfunction	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general sluggishness, loss of appetite, etc.
Kidney damage	Fever, rash, decreased urine output, general swelling, general sluggishness, joint pain, diarrhea, etc.
Interstitial pneumonia	Shortness of breath or difficulty breathing on exertion, such as when climbing stairs, dry cough and fever, etc. These symptoms may appear suddenly or persist.
Asthma	Difficulty breathing or a wheezing or whistling sound when breathing.

3 The following symptoms may appear after taking this medicine. If these symptoms persist or worsen, discontinue use and consult with your doctor, pharmacist or registered seller, bringing this instruction sheet with you

Drowsiness

4 If symptoms do not improve after taking this medicine 5 to 6 times, discontinue use and consult with your doctor or pharmacist, bringing this instruction sheet with you

Efficacy

◆ Relief from headache, toothache, pain after tooth extraction, menstrual pain (period pain), nerve pain, muscle pain, lower back pain, pain due to stiff shoulders, joint pain, sore throat (throat pain), ear ache, pain associated with bruising, pain associated with bone fractures, pain associated with sprains, and traumatic pain

◆ Fever relief during chills (feeling cold due to fever) and fevers

Headache

Fever

Toothache

Usage/Dosage

Take the following dosage with room-temperature or lukewarm water. Avoid taking on an empty stomach as much as possible. Wait at least 4 hours between doses.

Age	15 years of age or older	11 to 14 years of age	7 to 10 years of age	3 to 6 years of age	1 to 2 years of age	Under 1 years of age
Single dose	1 sachet	2/3 sachet	1/2 sachet	1/3 sachet	1/4 sachet	Do not take this medicine
Number of doses	Up to 3 times a day					

Note:

(1) Strictly follow the prescribed usage and dosage.

(2) If this medicine is being taken by a child, ensure it is used under the guidance and supervision of a parent or guardian.

Ingredients

In each sachet (1.6 g)

Components	Amount	Action
Acetaminophen	265 mg	Relieves fever and pain such as headache, toothache and menstrual pain.
Ethenzamide	300 mg	Analgesic effect to relieve pain.
Bromvalerylurea	200 mg	Sedative effect helps to relieve pain through acetaminophen, etc.
Anhydrous caffeine	50 mg	Relieves headache.

Additives: Lactose, carmellose calcium, hydroxypropyl cellulose, potato starch, magnesium aluminometasilicate, *l*-menthol

Precautions for Storage and Handling

(1) Store in a cool, dry place that is out of direct sunlight.
(2) Keep out of reach of children.
(3) Do not transfer to another container. (This may result in misuse or a change in quality)
(4) If you are dividing the contents of a sachet into multiple doses, store with the opening folded over and take the remainder of the medicine within 2 days.
(5) Do not take this medicine after it is past its expiration date.

Avoid taking on an empty stomach

☆ Ingredients that generally suppress fever and pain can irritate the stomach. Where possible, eat and drink something before taking this drug.

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.