

NARON ACE PLUS

- ◆ NARON ACE PLUS is an antipyretic analgesic containing two antipyretic and analgesic components combined with a gastric mucosal protective component.
- ◆ Bromovalerylurea and anhydrous caffeine assist the analgesic effects of Ibuprofen and Ethenzamide that block the cause and transmission, respectively, of pain, relieving pain/fever symptoms.
- ◆ The dried aluminum hydroxide gel protects gastric mucosa and prevents damage to the stomach.
- ◆ A small, easy-to-swallow tablet.

Indication

- ◆ Relief of headache, painful menses (menstrual pain), toothache, pain after tooth extraction, lumbago, pain due to stiff shoulders, muscular pain, joint pain, contusion pain, pain associated with sprain (sprain pain), bone fracture pain, traumatic pain, neuralgia, sore throat (throat pain) and earache
- ◆ Relief of fever during chills (feeling cold due to fever) and fever

Dosage and administration

Take the following dose with cold or lukewarm water. Avoid taking the drug on an empty stomach whenever possible. Provide a dosing interval for at least 4 hours.

15 years or over: 2 tablets per dose, 3 times daily

Under 15years: Do not take

Precaution

- Comply with the prescribed dosage and administration instructions.

How to take out the tablets

- Press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets. (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa)

Ingredient

In 2 tablets

Ibuprofen 144mg

Ethenzamide 84mg

Bromvalerylurea 200mg

Anhydrous caffeine 50mg

Dried aluminum hydroxide gel 66.7mg (200mg in 6 tablets)

Excipients

Anhydrous silicic acid, hydroxypropylcellulose, cellulose, hypromellose, croscarmellose sodium, magnesium stearate, lactose, talc, brilliant blue

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur)

- This product should not be taken in the following persons
 - Patients who have had an allergic symptom to this drug or its ingredients.
 - Patients who have experienced asthma from taking this drug, other antipyretic analgesics, cold remedies.
 - Children under 15 years old.
 - Pregnant women who are expected to give birth within 12 weeks.
- This medicine should not be taken together with any of the following medicines
 - Other antipyretics/analgesics, cold medicine, sedatives, medicine for motion sickness
- After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness may occur)
- Do not drink alcohol before and after taking this drug
- Do not take this medicine for a long time

Consultation

- The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before administration
 - Patients undergoing medical treatment from a physician or dentist.
 - Pregnant women or women suspected of being pregnant.
 - Nursing women.
 - The elderly.
 - Patients who have experienced allergic symptoms associated with drugs, etc.
 - Persons diagnosed as having the following.
 - Heart disease, kidney disease, liver disease, systemic lupus erythematosus, mixed connective tissue disease
 - Patients who experienced any of the following diseases.
 - Gastric/duodenal ulcer, ulcerative colitis, Crohn's disease
- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, dentist, pharmacist, or registered salesperson for a consultation
 - Skin: rash/redness, itching, and bruises
 - Gastrointestinal system: nausea/vomiting, loss of appetite, gastric discomfort, stomach pain, stomatitis, heartburn, heaviness in the stomach, gastrointestinal bleeding, abdominal pain, diarrhea, and bloody stool
 - Neuropsychiatric system: dizziness
 - Cardiovascular system: palpitations
 - Respiratory system: shortness of breath
 - Other: blurred vision, tinnitus, oedema, nosebleeds, bleeding from the gums, defects in hemostasis, bleeding, back pain, excessive drop of body temperature, and general malaise
- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.
 - Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.
 - Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, etc. may persist or suddenly worsen.
 - Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.
 - Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.
 - Aseptic meningitis: Symptoms, such as severe headache with a tight sensation along the neck, fever and feeling nausea/vomiting, etc. may occur. (these symptoms are frequently reported particularly in patients under treatment for systemic lupus erythematosus or mixed connective tissue disease.)
 - Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.
 - Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.
 - Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
 - Constipation, drowsiness
- If the symptoms do not resolve after taking 5 to 6 doses, discontinue the use of this drug, and bring this document to a physician, dentist, pharmacist, or registered salesperson for a consultation

Precautions for storage and handling

- Avoid direct sunlight, and store the product in a cool place with little humidity.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not take the product past the expiration date.

11

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.