

ARCHSKIN

Newly designated quasi-
drugs

Bactericidal disinfectants
(including special adhesive
plasters)

◆ARCHSKIN is a simple and convenient liquid bandage.■When it is spread on wound area, it forms a bactericidal clear coating and protects the wound from dusts and germs.■It is not easily removed and protects wound at situations involving the use of water, such as while cooking, washing, and bathing in the bath or in sea water.

Indication

Disinfection and protection (covering) of cuts, abrasion, stab wounds, scratch, shoe sores, and wound surface

Dosage and administration

Please sufficiently wipe off water or substances adhering to the wound surface, and apply appropriate amount on the wound and dry it.

Comply with the prescribed dosage and administration instructions.

Use for children should be supervised by a parent.

Attention should be paid so that the medicine will not come into contact with eyes. In case the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water, and immediately seek an ophthalmologist for medical treatment.

This is for external use only.

It might irritate the wound for a moment; however, it quickly creates a coating and protects the wound.

If the wound area is wide, put a small amount of absorbent cotton or gauze on the wound and apply the liquid drug over it.

Although this causes a slight pain, this is the most stable and effective approach.

If you would prefer to quickly remove the coating of wound area, spread Archskin on the coating and wipe it off with paper or cloth while slowly dissolving the coating.

ingredient and amount

In 100 g
Trichlorocarbanilide 0.1g

Excipients

celluloid, Isopropyl myristate, Anhydrous ethanol, Ethyl acetate,,

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

Do not use in the following sites.

sores
erosion

The purulent lesion

Consultation

The following persons should contact a physician or pharmacist for a consultation before using the drug:

Persons who or whose family members have an allergic predisposition

Persons who have had an allergic symptom to drugs

In the following cases, immediately discontinue this drug, and consult a physician or pharmacist. Be sure to show this leaflet.

The following symptoms may occur after using the drug:

Skin

rash/redness

itching

For instances wherein symptoms are not relieved despite taking this medicine

Precautions for storage and handling

Store in a cool, low humidity place away from direct sunlight.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not place the drug close to any source of flame.

Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.