

SAMON ACE

Class 2 OTC drug

Nutritional supplements

◆Physical dullness, stiff shoulders, lower back pain, and numbness in limbs are symptoms often seen as you get old. ◆SAMON ACE is a preparation that contains Japanese and Chinese drug extracts as main ingredients, as well as ginseng extract and vitamin E, and effective for these symptoms associated with aging.

Indication

Following symptoms observed in middle age and older: malaise, stiff shoulders, lower back pain, thirst, coldness/numbness in limbs, pollakiuria

Dosage and administration

Adults: Take 1 capsule per dose 3 times daily with cold or lukewarm water.

Comply with the prescribed dosage and administration instructions.

•How to take out the capsules

•Press the convex part of the PTP sheet containing the capsule firmly with your fingertips, break the aluminum foil on the back side, and take out the capsule. (If the capsules are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 3 capsules

Chibakujiogan dry extract 866mg (equivalent to 2000mg of Rehmannia root, 1000mg of Dioscorea rhizome, 1000mg of Cornus fruit, 1000mg of Moutan bark, 1000mg of Alisma rhizome, 1000mg of Poria sclerotium, 1000mg of Anemarrhena rhizome, 666mg of Phellodendron bark)

Ginseng extract 70mg (equivalent to 1000 mg of the crude drug)

Tocopherol Calcium Succinate 10mg

Excipients

Aluminum hydroxide/magnesium hydroxide, cellulose, Magnesium stearate, Brilliant blue, Erythrosine, Sunset Yellow FCF, Sodium lauryl sulfate, Gelatin

Precautions

Consultation

●If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Gastrointestinal system

Loss of appetite

●The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.
soft stool

●When symptoms do not improve even after taking this medicine for a while, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not take the product past the expiration date.

Even before the expiration date, use up to within 6 months after it is opened (to assure quality).

12

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.