

Make sure to read this instruction sheet before taking this medicine.  
Additionally, store this sheet so that it can be referenced when needed.

# ACETAMINOPHEN TABLETS s

Contains the same active ingredients as medical drugs\*

Product Name: ACETAMINOPHEN TABLETS s  
<Antipyretic analgesic>

Can be taken from age 5

Class 2 OTC drug

- ◆ ACETAMINOPHEN TABLETS s is an antipyretic analgesic containing acetaminophen, which relieves fever, headache, menstrual pain, and other types of pain.
  - ◆ Can be taken by the whole family from age 5 and up.
  - ◆ Contains the same active ingredients as medical drugs\*.
- Note: Dosage differs from that of prescription drugs.



## Precautions for Use



### Things You Must Not Do



(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects or accidents occurring)

- The following persons should not take this medicine:
  - Persons who have had allergic symptoms in the past as a result of using this product or any of its components.
  - Persons who have experienced asthma while taking this medicine or other antipyretic analgesics or cold medicines.
- Do not take any of the following medicines while taking this medicine:
  - Other antipyretic analgesics, cold medicines, sedatives
- Do not drink alcohol before or after taking this medicine
- Do not use for long periods of time



## Things To Consult On



- The following individuals should consult with their doctor, dentist, pharmacist, or a registered seller before taking this medicine:
  - Persons receiving treatment from a doctor or dentist.
  - Persons who are pregnant or who think they may be pregnant.
  - Elderly persons.
  - Persons who have had allergic symptoms due to medication or similar substances in the past.
  - Persons who have received the following diagnoses:
    - Heart disease, kidney disease, liver disease, stomach and duodenal ulcers
- If any of the following symptoms appear after you have taken this medicine, you may be experiencing side effects. As such, discontinue use immediately and consult with your doctor, pharmacist, or registered seller, bringing this instruction sheet with you

Relevant Area	Symptoms
Skin	Rash, redness, itching
Digestive system	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Other	Hypothermia

In rare cases, the serious side effects listed on the right may occur.  
In such cases, see a doctor immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Itching, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion immediately after taking this medicine.
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	High fever, bloodshot eyes, eye discharge, lip sores, sore throat, widespread skin rash/redness, small bumps (small pustules) appearing on reddened skin, full-body sluggishness, loss of appetite, etc., that persists or worsens rapidly.
Drug-induced hypersensitivity syndrome	Symptoms include widespread reddening of the skin, a rash over the entire body, fever, bodily sluggishness, and swelling of the lymph nodes (in the neck, armpits, groin, etc.).
Liver dysfunction	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general sluggishness, loss of appetite, etc.
Kidney damage	Fever, rash, reduced urine output, whole-body swelling, general sluggishness, joint pain (joints are sore), diarrhea
Interstitial pneumonia	Shortness of breath or difficulty breathing on exertion, such as when climbing stairs, dry cough and fever, etc. These symptoms may appear suddenly or persist.
Asthma	Difficulty breathing or a wheezing or whistling sound when breathing.


- If symptoms do not improve after using this medicine 5 to 6 times, discontinue use and consult with your doctor, dentist, pharmacist or registered seller, bringing this instruction sheet with you

## Efficacy

- ◆ Relief from headache, painful menses (menstrual pain), toothache, pain after tooth extraction, lower back pain, pain due to stiff shoulders, muscle pain, joint pain, pain associated with bruising, pain associated with sprains, pain associated with bone fractures, traumatic pain, nerve pain, sore throat (throat pain) and ear ache
- ◆ Fever relief during chills (feeling cold due to fever) and fevers



Headache



Menstrual pain






Fever



Toothache

## Usage/Dosage

Take the following dosage with room-temperature or lukewarm water. Avoid taking on an empty stomach as much as possible. Wait at least 4 hours between doses.

Age	Single dose	Number of doses
Adults (15 years and older)	3 tablets 	Up to 3 times a day
11 to 14 years of age	2 tablets 	
5 to 10 years of age	1 tablet 	
Under 5 years old	Do not take this medicine	

Note:

- (1) Strictly follow the prescribed usage and dosage.
- (2) If this medicine is being taken by a child, ensure it is used under the guidance and supervision of a parent or guardian.

(3) How to take out the tablets

As shown in the illustration, press the convex part of the PTP sheet containing the tablet hard with a fingertip to tear the aluminum foil on the reverse side, remove the tablet, and take it. (Accidental ingestion of the PTP sheet as-is may lead to unexpected accidents, such as penetration into the esophageal mucosa.)

<Tablet removal diagram>

Push it out



The medicine should be taken as follows:

Take as soon as the pain starts.



For adults, take 3 tablets each time with a cup of room-temperature or lukewarm water.



As much as possible, eat or drink beforehand to avoid taking the medicine on an empty stomach.



## Ingredients

In 9 tablets

Ingredients	Amount	Action
Acetaminophen	900 mg	Acts on the central nervous system to relieve fever, headaches, menstrual pain, and other types of pain.

Additives: Anhydrous silicic acid, cellulose, crosslinked sodium carboxymethylcellulose (CMC-Na), hydroxypropyl cellulose, hydrogenated oil, magnesium stearate

## Precautions for Storage and Handling

- (1) Store in a cool, dry place that is out of direct sunlight.
- (2) Store this product out of reach of children.
- (3) Do not transfer to another container. (This may result in misuse or a change in quality)
- (4) Do not take the product after the expiration date. Use this medicine within 6 months of opening, even if still within the expiration date. (In order to ensure quality.)



### Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.