

PABRON RHINITIS CAPSULES Sa

Designated Class 2 OTC
drug

Medicine for rhinitis

◆PABRON RHINITIS CAPSULES Sa is an oral drug used to treat rhinitis. Twice daily doses achieve an excellent effect. ■ The white granules dissolve quickly and begin to work, while the orange granules dissolve gradually and effectively relieve annoying rhinitis symptoms after a while.

Indication

Relief of the following symptoms due to acute rhinitis, allergic rhinitis or sinusitis; sneezing, runny nose (excessive nasal discharge), stuffy nose, watery eyes, sore throat, dull headache (heaviness in the head)

Dosage and administration

Take the following dose with cold or lukewarm water every 12 hours.

15 years or over: 2 capsules per dose, twice daily

Under 15 years: Do not take

Comply with the prescribed dosage and administration instructions.

•How to take out the capsules

•Press the convex part of the PTP sheet containing the capsule firmly with your fingertips, break the aluminum foil on the back side, and take out the capsule. (If the capsules are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 2 capsules

Pseudoephedrine hydrochloride 60mg

Carbinoxamine maleate 6mg

Belladonna total alkaloid 0.2mg

Anhydrous caffeine 50mg

Excipients

Magnesium aluminometasilicate ,Sugar,Corn starch ,Hydroxypropylcellulose,Magnesium stearate,cellulose,Talc,Ammonio alkyl methacrylate copolymer,Stearyl alcohol ,Sorbitan trioleate,Sunset Yellow FCF ,Gelatin,Sodium lauryl sulfate

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Persons who have had an allergic symptom to this drug or its ingredients.

Persons with the following symptoms:

Urination difficulty due to prostatic hypertrophy

Persons diagnosed as having the following

thyroid disease

heart disease

Hypertension

Diabetes

This drug should not be taken together with the following drugs:

Other internal medicine for rhinitis, internal medicine containing antihistamines (including cold medicine, expectorant, medicine for motion sickness, medicine for allergies), indigestion and heartburn relief.

Oral drugs that contain pseudoephedrine hydrochloride or pseudoephedrine sulfate.

●After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur).

● Do not take this medicine for a long time

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

Nursing women.

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

Patients who have experienced insomnia, dizziness, lassitude, shivering, palpitations from taking cold remedies, antitussives and expectorants, oral remedies for rhinitis, etc.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

Glaucoma

kidney disease

Patients who are being treated with a monoamine oxidase inhibitor (selegiline hydrochloride, etc.)

(Selegiline hydrochloride is used to treat Parkinson's disease.)

●If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Gastrointestinal system

Vomiting

Vomiting

Loss of appetite

Neuropsychiatric system

dizziness

Insomnia

Hypersensitivity

headache

Convulsions

Urinary system:

urination difficulty

Other:

Facial hot flashes

abnormal brightness

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Acute generalised exanthematous pustulosis Some symptoms, such as hyperthermia, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, may persist or suddenly worsen.

●The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Thirst, sleepiness, constipation, blurred vision

●When symptoms do n...

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened (to assure the quality).

21

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.