

# PABRON S GRANULES

- ◆ PABRON S GRANULES are a cold remedy with an effect on 11 cold symptoms such as sputum and a sore throat, containing 7 active ingredients including bromhexine hydrochloride, an ingredient caring for the respiratory tract mucosa and vitamin C (contained in the form of calcium ascorbate).
- ◆ They are non-caffeinated and easy to take, with a citrus flavor.
- ◆ Since they can be taken by children of 1 year and older, they are suitable for use as part of your stock of household medications.

## Indication

To alleviate cold symptoms (cough, sputum, sore throat, sneezing, runny nose, stuffy nose, chills (chill due to fever), fever, headache, joint pain, and muscle pain)

## Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

Adults (15 years or over): 1 packet per dose, 3 times daily

11 to 14 years: 2/3 packet per dose, 3 times daily

7 to 10 years: 1/2 packet per dose, 3 times daily

3 to 6 years: 1/3 packet per dose, 3 times daily

1 to 2 years: 1/4 packet per dose, 3 times daily

Under 1 year: Do not take

## Precaution

- Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.
- Infants under 2 years old should seek medical care from a physician first. Use the drug only in an unavoidable case.

## Ingredient

In 1 packet (0.96g)

Bromhexine hydrochloride 4mg

Dextromethorphan Hydrobromide 16mg

dl-Methylephedrine Hydrochloride 20mg

Acetaminophen 300mg

Chlorpheniramine Maleate 2.5mg

Calcium ascorbate (Calcium ascorbate) 83.3mg (250mg in 3 packets)

Riboflavin (vitamin B<sub>2</sub>) 4mg

## Excipients

Potato starch, anhydrous silicic acid, cellulose, D-mannitol, hydroxypropylcellulose, aspartame (L-phenylalanine compound), flavoring, starch sodium octenyl succinate

## Precaution

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B<sub>2</sub> contained in the drug.

## Precautions

### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur)

- This product should not be taken in the following persons
  - Patients who have had an allergic symptom to this drug or its ingredients.
  - Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.
- Do not use any of the following medications while taking this drug
  - Other cold medicines, antipyretics/analgesics, sedatives, expectorants, and internal medicines, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.)
- After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness may occur)

- Do not drink alcohol before and after taking this drug
- Do not take this medicine for a long time

### Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Nursing women.

The elderly.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms.

Hyperthermia, urination difficulty

Persons diagnosed as having the following.

Thyroid disease, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Skin: rash/redness, itching

Gastrointestinal system: nausea/vomiting, and loss of appetite

Neuropsychiatric system: dizziness

Respiratory system: shortness of breath, breathing difficulties

Urinary system: urination difficulty

Other: excessive temperature decrease

- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Drug-induced hypersensitivity syndrome: Symptoms such as widespread redness of the skin, generalised rash, fever, malaise, and swelling of lymph nodes (neck, armpit, groin, etc.) may occur.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.

Interstitial pneumonia: Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance.

Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.

Aplastic anemia: symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.

Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Dry mouth, drowsiness

- When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you

### Precautions for storage and handling

- Avoid direct sunlight, and store the product in a cool place with little humidity.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.
- Do not take the product past the expiration date.

**【Disclaimer on Multilingual Product Information】**

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.