

# PABRON S GOLD W TABLETS

Designated Class 2 OTC  
drug

Cold Remedies

◆PABRON S GOLD W TABLETS are a compound of 6 active ingredients including ambroxol hydrochloride and L-carbocysteine. A cold medication that is effective on 11 symptoms associated with the common cold, such as sore throat, cough, and runny nose. It is suitable for family use.

## Indication

Relief of various symptoms of a common cold: running nose, stuffy nose, sneezing, sore throat, cough, phlegm (sputum), chills (feeling cold due to fever), fever, headache, joint pain, and muscle pain.

## Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

15 years or over: 2 tablets per dose, 3 times daily

Children aged 12–14 years old: 1 tablet, 3 times a day

Children under 12 years old: Do not take this drug.

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

Do not return the tablets touched by wet fingers into the jar. (It will cause discoloration and may deteriorate the quality.)

## ingredient and amount

In 2 tablets

Ambroxol hydrochloride 15mg

L-carbocysteine 250mg

Dihydrocodeine Phosphate 8mg

Acetaminophen 300mg

Chlorpheniramine Maleate 2.5mg

Riboflavin 4mg

## Excipients

cellulose, anhydrous silicic acid, calcium hydrogen phosphate, Sodium starch glycolate, Hydroxypropylcellulose, hardened oil, Magnesium stearate

## Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

## Precautions

### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Persons who have had an allergic symptom to this drug or its ingredients.

Children under 12 years old.

Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.

This drug should not be taken together with the following drugs:

Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.) and gastrointestinal analgesics

● After taking this drug, do not drive a car or operate machinery (sleepiness may occur).

For those who are lactating, do not use this medicine or avoid lactating while using this medicine.

Do not drink alcohol before and after taking this drug.

● Do not take this medicine for a long time

## Consultation

● The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

heart disease

Liver disease

kidney disease

Gastric/duodenal ulcer

Glaucoma

Respiratory functional disorder, obstructive sleep apnea syndrome, adiposity

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Gastrointestinal system: nausea/vomiting, loss of appetite, feeling of discomfort in the stomach, stomach ache, abdominal pain, feeling of stomach/abdominal distention, heartburn, and diarrhea

Neuropsychiatric system

dizziness

Numbness

Urinary system:

urination difficulty

Other:

oedema

Excessive temperature decrease

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised

exanthematous pustulosis: Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), ...

## Precautions for storage and handling

Store in a cool, low humidity place away from direct sunlight.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened (to assure the quality).

41

### 【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.