

PABRON KIDS COLD TABLETS

Class 2 OTC drug

Cold Remedies

◆PABRON KIDS COLD TABLETS is a cold medication for children that has steady effect on cough, nasal and throat symptoms, and fever. □
◆The sweet small tablets are designed for kids -i.e., easy to swallow, and does not contain caffeine that disturbs sleep. ◆Acute cough, fever, and runny nose should be relieved at any cost. Treat with Pabron Kids Cold tablets at an early stage, keep the body warm, and have a good rest.

Indication

To alleviate cold symptoms (sore throat, fever, cough, sputum, runny nose, stuffy nose, sneezing, chill (chill due to fever), headache, joint pain, muscle pain)

Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

11 to 14 years: 4 tablets per dose, 3 times daily

7 to 10 years: 3 tablets per dose, 3 times daily

5 to 6 years: 2 tablets per dose, 3 times daily

Under 5 years: Do not take

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

How to take out the tablets

•As shown in the figure, press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets.

• (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 4 tablets

Tipepidine Hibenazate 16.7mg

Guaifenesin 33.3mg

Chlorpheniramine Maleate 1.6mg

Acetaminophen 200mg

Excipients

cellulose,anhydrous silicic acid ,Magnesium aluminometasilicate ,Lactose ,Carboxymethylcellulose

calcium ,Hydroxypropylcellulose,Magnesium stearate,Hypromellose,Sugar,Titanium oxide,Ferric oxide ,Carnauba wax ,flavor,Vanillin,Benzyl benzoate

Precautions

When not to use the product

Although the drug is intended for children, the document describes the general precaution for cold medicine (If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.

● This medicine should not be taken together with any of the following medicines:

Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.) and gastrointestinal analgesics

●After taking this drug, do not drive a car or operate machinery (sleepiness may occur).

Do not drink alcohol before and after taking this drug.

● Do not take this medicine for a long time

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

heart disease

Liver disease

kidney disease

Gastric/duodenal ulcer

Glaucoma

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Gastrointestinal system

Vomiting

Vomiting

Loss of appetite

Neuropsychiatric system

dizziness

Urinary system:

urination difficulty

Other:

Excessive temperature decrease

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur...

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not take the product past the expiration date.

21

【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.