

PABRON KIDS COLD GRANULES

Class 2 OTC drug

Cold Remedies

◆PABRON KIDS COLD GRANULES is a cold medication for children that has steady effect on cough, nasal and throat symptoms, and fever. ■ The particles are designed for kids -i.e., flavored with strawberries that kids love for easy administration, and does not contain caffeine that disturbs sleep. ■ Acute cough, fever, and runny nose should be relieved at any cost. Treat with Pabron Kids Cold Particle at an early stage, keep the body warm, and have a good rest.

Indication

To alleviate cold symptoms (sore throat, fever, cough, sputum, runny nose, stuffy nose, sneezing, chill (chill due to fever), headache, joint pain, muscle pain)

Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

7 to 10 years: 1 packet per dose, 3 times daily

3 to 6 years: 2/3 packet per dose, 3 times daily

1 to 2 years: 1/2 packet per dose, 3 times daily

Under 1 year: Do not take

Comply with the prescribed dosage and administration instructions.

Use for children should be supervised by a parent.

Infants under 2 years old should seek medical care from a physician first. Use the drug only in an unavoidable case.

ingredient and amount

In 1 packet (0.96 g)

Tipepidine Hibenzone 12.5mg

Guaifenesin 25mg

Chlorpheniramine Maleate 1.2mg

Acetaminophen 150mg

Excipients

anhydrous silicic acid ,Corn starch ,Sugar,D-Mannitol ,Aspartame (L- phenylalanine compound),Acesulfame K,Sucralose ,Hydroxypropylcellulose,Erythrosine,Magnesium stearate,flavor,Vanillin,Ethyl vanillin

Precautions

When not to use the product

Although the drug is intended for children, the document describes the general precaution for cold medicine (If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.

Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.) and gastrointestinal analgesics

● This medicine should not be taken together with any of the following medicines:

● After taking this drug, do not drive a car or operate machinery (sleepiness may occur).

Do not drink alcohol before and after taking this drug.

● Do not take this medicine for a long time

Consultation

● The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

heart disease

Liver disease

kidney disease

Gastric/duodenal ulcer

Glaucoma

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a

consultation.
Skin
rash/redness
itching
Gastrointestinal system
Vomiting
Vomiting
Loss of appetite
Neuropsychiatric system
dizziness
Urinary system:
urination difficulty
Other:
Excessive temperature decrease

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.
Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur...

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.

Do not take the product past the expiration date.

21

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.