

PABRON RHINITIS CHEWABLE TABLETS EX

Designated Class 2 OTC drug

Medicine for rhinitis

◆ PABRON RHINITIS CHEWABLE TABLETS EX is an oral anti-rhinitis drug that quickly dissolves in the mouth to exert an excellent effect on acute rhinitis symptoms.■ It can be taken without water and easily at any place.■ It has a milk and menthol flavor giving a cooling sensation.

Indication

Relief of the following symptoms due to acute rhinitis, allergic rhinitis or sinusitis; sneezing, runny nose (excessive nasal discharge), stuffy nose, watery eyes, sore throat, dull headache (heaviness in the head)

Dosage and administration

Take the following dose by chewing or dissolving the drug in the mouth. However, provide a dosing interval for at least 4 hours.

15 years or over: 2 tablets per dose, 3 times daily

7 to 14 years: 1 tablet per dose, 3 times daily

Under 7 years: Do not take

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

How to take out the tablets

•As shown in the figure, press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets.

• (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 2 tablets

Pseudoephedrine hydrochloride 40mg

dI-Methylephedrine Hydrochloride 10mg

Chlorpheniramine Maleate 2mg

Dipotassium Glycyrrhizinate 15mg

Belladonna total alkaloid 0.2mg

Anhydrous caffeine 25mg

Excipients

D-Mannitol ,Powdered sugar,Corn starch ,cellulose,anhydrous silicic acid ,Crosopovidone,Hydroxypropylcellulose,Aspartame (L- phenylalanine compound),Sucralose ,Calcium stearate,l-Menthol,flavor,,

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Persons who have had an allergic symptom to this drug or its ingredients.

Urination difficulty due to prostatic hypertrophy

Persons diagnosed as having the following

thyroid disease

heart disease

Hypertension

Diabetes

This drug should not be taken together with the following drugs:

Other internal medicine for rhinitis, internal medicine containing antihistamines (including cold medicine, expectorant, medicine for motion sickness, medicine for allergies), indigestion and heartburn relief.

●After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur).

● Do not take this medicine for a long time

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

Nursing women.

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

Patients who have experienced insomnia, dizziness, lassitude, shivering, palpitations from taking cold remedies, antitussives and expectorants, oral remedies for rhinitis, etc.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

Glaucoma

kidney disease

Patients who are being treated with a monoamine oxidase inhibitor (selegiline hydrochloride, etc.)

(Selegiline hydrochloride is used to treat Parkinson's disease.)

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Gastrointestinal system

Vomiting

Vomiting

Loss of appetite

Neuropsychiatric system

dizziness

Insomnia

Hypersensitivity

headache

Convulsions

Urinary system:

urination difficulty

Other:

Facial hot flashes

abnormal brightness

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Acute generalised exanthematous pustulosis Some symptoms, such as hyperthermia, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, may persist or suddenly worsen.

Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of sk...

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened (to assure the quality).

11

【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.