

PABRON SUPPOSITORIES for KIDS

Class 2 OTC drug
antipyretics

◆Fever is the most common symptom as it is said that all the pediatric diseases start with fever. However, it is important to alleviate fever because it could consume physical strength. ■Children's pabron suppository effectively relieves acute fever in children.

Indication

To temporarily alleviate pediatric fever

Dosage and administration

Insert the following dose into the anus.

6 to 12 years: 1 to 2 pieces per dose, once daily

3 to 5 years: 1 piece per dose, once daily

1 to 2 years: 1/2 to 1 piece per dose, once daily

Under 1 year: Do not use

Comply with the prescribed dosage and administration instructions.

Use for children should be supervised by a parent.

Do not use the drug for 2 days in a row.

Use the drug only for the anus. Do not take it orally.

ingredient and amount

In 1 piece (1 g)

Acetaminophen 100mg

Excipients

Hard fat

Precautions

When not to use the product

Although the drug is intended for children, the document describes the general precaution for antipyretics (If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.

●This drug should not be taken together with the following drugs:

Other cold medicines, antipyretics/analgesics, sedatives

● Do not drink alcohol before/after using this medicine

● Do not take this medicine for a long time

Consultation

●The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before use.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

The elderly

Persons with a weak constitution

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons diagnosed as having the following

heart disease

Liver disease

kidney disease

Gastric/duodenal ulcer

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Gastrointestinal system

Vomiting

Vomiting

Loss of appetite

diarrhea

soft stool

Urge to defecate

Neuropsychiatric system

dizziness

Other:

Excessive temperature decrease

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after use.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.

Interstitial pneumonia: Shortness ...

Precautions for storage and handling

Avoid direct sunlight, and store in a cool place with little humidity and temperature below 30°C.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)

For storage, store in the outer box, place the tip downward, and stand them according to the instruction on the outer box.

Do not take the product past the expiration date.

Even within the expiration date, use up as soon as possible once it is opened (to assure the quality).

91

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.