

PRESER ACE INSERTING-TYPE OINTMENT T

- ◆PRESER ACE INSERTING-TYPE OINTMENT T contains active ingredients to relieve piles with annoying pain/acute bleeding such as lidocaine that relieves pain, tetrahydrozoline hydrochloride that inhibits bleeding, and hydrocortisone acetate that relieves inflammation, and achieves an excellent effect.
- ◆The drug has 2 usages - insert the drug for internal hemorrhoids (deep in the anus), and apply the drug for hemorrhoids outside the anus. You can insert the drug cleanly without directly touching the affected site and drug.
- ◆It soothes symptoms, and gives comfortable sense of use.

Indication

[Infusion] To alleviate pain/bleeding/swelling/itching due to bleeding piles/blind piles

[Application] To alleviate pain/bleeding/swelling/itching due to bleeding piles/blind piles and for disinfection

Dosage and administration

[Injection]

Insert the tip of container into the anus, and inject the entire amount.

15 years or over: 1 piece per dose, 1 to 3 times daily

Under 15 years: Do not use

- Remove the cap, and squeeze out a little amount of ointment to make the surface smooth.
- Insert the tip of the container into the anus, and press the container to inject the drug. (Pull off the tip while pressing the container.)

[Application]

Apply the following dose onto the anus. Do not use the drug for injection once it is applied on the lesion.

15 years or over: suitable dose, 1 to 3 times daily

Under 15 years: Do not use

- Remove the cap, apply the ointment directly, or spread on a gauze, and then put it on the affected site.
 - * If the ointment is hard and difficult to squeeze out, warm the container with palms to make the ointment soft.
 - * Even though a small amount of the drug remains in the injectable container, a proper dose has already been injected in the affected site.

Precaution

- Comply with the prescribed dosage and administration instructions.
- Do not use the drug for children.
- The medicine must be used only for the anal area.
- Insert only the tip of the container into the anus.

Ingredient

In 1 piece (2g)

Hydrocortisone acetate 5mg

Tetrahydrozoline hydrochloride 1mg

Lidocaine 60mg

I-Menthol 10mg

Allantoin 20mg

Tocopherol acetate 60mg

Chlorhexidine Hydrochloride 5mg

Excipients

Liquid paraffin, isopropyl myristate, white beeswax, carboxyvinyl polymer, olive oil, petrolatum

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- This product should not be used in the following persons:

Persons who have had an allergic symptom to the drug or the drug ingredient, namely, chlorhexidine.

Patients with purulent hemorrhoidal lesions.

- Do not take this medicine for a long time

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered sales person for a consultation.

Skin: rash/redness, itching, swelling

Other: Irritation, Purulence

- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after use.

- If the symptoms do not resolve after taking this drug for about 10 days, stop taking this medicine, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Precautions for storage and handling

- Avoid direct sunlight, and store the product in a cool place with little humidity.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)
- Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).
- Do not flush the container after using the product in the toilet.

31

[Disclaimer on Multilingual Product Information]

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.