

# RiUP X5 Plus Lotion

Category I Medicines

Hair growth agent for  
premature alopecia

The drug contains 5% minoxidil, an active ingredient, and promotes excellent hair growth. The container is designed to automatically apply a dose of 1mL when you open the cap, and press it to the scalp. In addition to minoxidil, the drug contains 3 active ingredients\* that freshen you up, and improve the environment for hair growth in the scalp. Pyridoxine hydrochloride, Tocopherol acetate, Menthol

## Indication

To promote hair growth in premature alopecia and prevent the progression of hair loss (fallen hair)

## Dosage and administration

For an adult man ( $\geq 20$  years old), apply 1 mL on the scalp with hair loss twice a day.

Use 1 mL per dose regardless of the size of the affected area. A dose of 1 mL is sufficient for spreading the drug on the entire scalp.

The container is designed to measure 1 mL.

A dose exceeding the range designated by the dosage and administration instructions or a more frequent application does not affect the efficacy. Strictly comply with the specified dosage and administration.

(If you use the drug with more than the prescribed dose, the effect hardly increases, and adverse reactions are more likely to occur.)

Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water. If the symptoms are severe, seek an ophthalmologist for medical treatment.

If you touch a mucous membrane (e.g., the eyes) with fingers stained with the drug solution, irritation will occur. Wash your hands well to remove the drug solution.

Do not stain objects that might melt with alcohol (glass frames, chemical fiber, etc.) with the drug.

Use hair styling agents and hair spray after applying the drug.

If you use hair dye (hair color, gray hair dye, etc.), apply the drug after the hair dye is completely finished.

## ingredient and amount

In 100 mL

Minoxidil 5.0g

Pyridoxine Hydrochloride (vitamin B6) 0.05g

Tocopherol Acetate 0.08g

l-Menthol 0.3g

## Excipients

Cetanol, 1,3-Butylene glycol, Butylated hydroxytoluene, phosphate, Anhydrous ethanol

## Precautions

### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Women

The safety of the drug has not been confirmed in Japanese women. Female patients should use RiUP series for women that contains 1% of minoxidil.

Minors (under 20 years old)

The product has not been used in Japan.

Persons with alopecia other than premature alopecia (e.g., alopecia areata and hair loss due to thyroid disease) or alopecia by an unknown cause

The drug is effective only for premature alopecia.

Acute hair loss and spotty hair loss

They are highly likely to be alopecia other than premature alopecia.

Do not use in the following sites.

Use the drug only for the scalp. Do not take it orally.

Blood pressure may decrease.

Scalp with wound, eczema or inflammation (redness), etc.

The drug may worsen the wound.

When you use this drug, do not use other hair growth agents and external preparations (ointment, liquid drug, etc.) in the scalp. When you use these other drugs, stop using this drug.

These other drugs may affect the absorption of this drug.

## Consultation

The following persons should contact a physician or pharmacist for a consultation before using the drug:  
Persons who have experienced allergic symptoms (e.g., rash/redness, itching, and eruption) associated with drugs and cosmetics.  
Persons with hypertension/ hypotension  
The drug may possibly affect blood pressure.  
Persons with heart or kidney disorder  
The drug may affect the heart and kidneys.  
Persons with oedema  
The drug may worsen oedema.  
Persons whose family and siblings have no premature alopecia.  
It is considered that genetic factors affect hugely the onset of premature alopecia.  
The elderly (≥65 years old)  
Generally, unfavorable symptoms are likely to occur in the elderly.  
Persons with the following diagnoses:  
Thyroid disease (hypothyroidism and hyperthyroidism)  
The hair loss may be caused by thyroid disease.  
If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to a physician or pharmacist for a consultation  
Skin: rash/redness in scalp\*, itching, eruption, dandruff, feeling hot in the application site, etc.  
Neuropsychiatric system: headache, faint, and dizziness  
Cardiovascular system: chest pain and increasing heart rate  
Metabolic system: sharp increase of body weight with unknown cause and oedema in the extremities  
\*: The symptoms may occur in sites other than the scalp.  
If none of the following improvements are achieved after using the drug for 6 months, stop using this medicine, and show this document to a physician or pharmacist for a consultation.  
Severity of hair loss, hair growth/soft hair growth, terminal hair growth, and amount of fallen hair (the decrease of not only bold fallen hair but also thin fallen hair is a sign of improvement).  
It may be alopecia other than premature alopecia, or the hair loss may be caused by other factors.  
If hair loss increases in severity or hair loss occurs in the condit...

## Precautions for storage and handling

After using the product, put the cap, avoid direct sunlight and high and extremely low temperatures, and store it in a cool place.  
Store the product beyond the reach of children.  
Do not transfer the drug to other containers to prevent misuse and quality deterioration.  
Do not place the drug close to any source of flame.  
Do not take the product past the expiration date.

14

### 【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.