

◆SEMPER QT is a quick-dissolving medication for motion sickness that can be taken without water. ■The product prevents and alleviates symptoms of motion sickness such as dizziness, nausea, and headache. ■Take the drug even after you begin to feel sick, and it will relieve dizziness, nausea, and headache.

Indication

1) prevention and relief of dizziness, nausea, and headache associated with motion sickness

Dosage and administration

Take the following dose by dissolving the drug in the mouth.

To prevent motion sickness, take 1 tablet per dose 30 minutes prior to boarding.

If you need to take an additional dose, then take 1 tablet per dose at an interval of at least 4 hours.

15 years or over: 1 tablet per dose, up to twice daily

Under 15 years: Do not take

Comply with the prescribed dosage and administration instructions.

How to take out the tablets

•Pull open the lid of the aluminum sheet that contains the tablets, pick the opening (red part) with your fingertips, and slowly peel it off. Press the tablet from the bottom and take out the tablets. (If the tablets are inadvertently taken together with the sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

Carefully remove the tablets from the package sheet to not break them.

Do not touch the tablets directly with wet hands. (Tablets easily dissolve in water.)

ingredient and amount

In 1 tablets

d-Chlorpheniramine Maleate 2mg

Scopolamine Hydrobromide 0.25mg

Excipients

Gelatin,D-Mannitol ,Aspartame (L- phenylalanine compound),flavor,Vitamin E

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This drug should not be taken together with the following drugs:

Other medicine for motion sickness, cold medicine, antipyretics/analgesics, sedatives, expectorant, indigestion and heartburn relief, internal medicine containing antihistamines, etc. (such as internal medicine for rhinitis and medicine for allergies)

●After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur).

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

urination difficulty

Persons diagnosed as having the following

Glaucoma

heart disease

●If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Neuropsychiatric system

headache

Urinary system:

urination difficulty

Other:

Facial hot flashes

abnormal brightness

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.

Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

●The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Thirst, sleepiness, constipation, blurred vision

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not take the product past the expiration date.

72

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.