

TOKUHON

Tokuhon is a plaster containing methyl salicylate that suppresses inflammation and relieves pain, vitamin E that improves blood circulation, and glycyrrhetic acid that alleviates skin inflammation, gentle for skin, and suitable for relieving muscle inflammation and pain. The four corners are rounded to prevent from being peeled off during use.

Indication

Stiff shoulders, lumbago, muscular pain, muscular fatigue, joint pain, contusions, sprains, frostbite

Dosage and administration

Peel off the paper liner on the drug surface, and apply the plaster onto the lesion. Repeat several times daily.

Precaution

- The use of the drug for children should be supervised by a parent. Children under 7 years old should not use this drug because it may cause tingling sensations.
- Clean the affected site before using the drug.
- If your skin is sensitive, do not apply the drug to the same site continuously.
- Put a small piece of the drug on the inner arm before use, and confirm that no rash/redness, itching, and eruption occurred.

Ingredient and amount

Active ingredients in 100g (0.54m²) of ointment

l-Menthol 6.5g
dl-Camphor 1.1g
Methyl Salicylate 7.3g
Vitamin E acetate 1.35g
Glycyrrhetic Acid 0.18g

Excipients

Zinc oxide, dibutyl hydroxytoluene, raw rubber, polyisoprene, ester gum, other two-component

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur)

- The medicine should not be used on the following parts
Periorbital area, mucosa, etc.
Eczema, contact dermatitis, open wounds

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use
Patients who have experienced allergic symptoms associated with drugs, etc.
- If you experience any symptoms such as rash/redness, itching, or eruption after use, these may be adverse reactions, so immediately discontinue the use, and consult with your physician, pharmacist, or registered salesperson with this explanatory document
- If symptoms do not resolve after using the drug for 5 or 6 days, stop using this medicine, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Precautions for storage and handling

- Store this in a cool place, avoiding direct sunlight.
- Store the product beyond the reach of children.
- After opening, use as soon as possible.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not use the product past the expiration date.

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.