

TOKUHON E

Class 3 OTC drug

Pain relief/antipruritic/astringent/anti-inflammatory (including cataplasm)

TOKUHON E is a scented external antiphlogistic analgesic agent containing glycol salicylate that suppresses inflammation and relieves pain, l-Menthol, and vitamin E that improves blood circulation. Since skin irritant contained in natural rubber (base composition) is removed by enzymes, the plaster causes less eruption and is gentle for skin. The four corners are rounded to prevent from being peeled off during use. The mocha pink color is less visible through clothes.

Indication

Muscular fatigue
joint pain, lumbago, muscular pain, pain due to stiff shoulders, contusion pain, bone fracture pain, pain associated with sprain (sprain pain), frostbite,

Dosage and administration

Peel the paper liner on the drug surface, and adhere it onto the lesion several times daily.
The use of the drug for children should be supervised by a parent. Children under 7 years old should not use this drug because it may cause tingling sensations.
Clean the affected site before using the drug.
If your skin is sensitive, do not apply the drug to the same site continuously.
Put a small piece of the drug on the inner arm before use, and confirm that no rash/redness, itching, and eruption occurred.

ingredient and amount

Active ingredients in 1 m²
l-Menthol 11.5g
Tocopherol acetate 2.5g
Glycol Salicylate 3g

Excipients

Zinc oxide, Butylated hydroxytoluene, Raw rubber, Ester gum, Other one-component

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)
Do not use in the following sites.
Periorbital area, mucosa, etc.
Eczema, contact dermatitis, open wounds.

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use. Patients who have experienced allergic symptoms associated with drugs, etc.
If you have symptoms such as rash/redness, itching, and eruption after use, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered sales person for a consultation.
- If symptoms do not improve even after using this medicine for 5 to 6 days, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.
Store the product beyond the reach of children.
After opening, fold up the opening of the bag, and use it as soon as possible.
To prevent misuse and quality deterioration, do not transfer the drug to container other than the attached storage bag.
Do not take the product past the expiration date.

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.