

TOKUHON E

Tokuhon E is a slightly scented topical antiphlogistic analgesic agent containing glycol salicylate, which suppresses inflammation and relieves pain, l-Menthol, and vitamin E, which improves blood circulation. This product is gentle to the skin, with rounded corners to resist peeling during use, and less visible through clothes thanks to its mocha pink color. The adhesive is not easily noticeable by others.

Indication

Stiff shoulders, lumbago, muscular pain, muscular fatigue, joint pain, contusions, sprains, frostbite

Dosage and administration

Peel off the paper liner on the drug surface, and apply the plaster onto the lesion. Repeat several times daily.

Precaution

- The use of the drug for children should be supervised by a parent. Children under 7 years old should not use this drug because it may cause tingling sensations.
- Clean the affected site before using the drug.
- If your skin is sensitive, do not apply the drug to the same site continuously.
- Put a small piece of the drug on the inner arm before use, and confirm that no rash/redness, itching, and eruption occurred.

Ingredient and amount

Active ingredients in 1m²

l-Menthol 11.5g

Vitamin E acetate 2.5g

Glycol salicylate 3.0g

Excipients

Zinc oxide, dibutyl hydroxytoluene, raw rubber, ester gum, other one-component

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur)

- The medicine should not be used on the following parts
Periorbital area, mucosa, etc.
Eczema, contact dermatitis, open wounds

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use
Patients who have experienced allergic symptoms associated with drugs, etc.
- If you experience any symptoms such as rash/redness, itching, or eruption after use, these may be adverse reactions, so immediately discontinue the use, and consult with your physician, pharmacist, or registered salesperson with this explanatory document
- If symptoms do not resolve after using the drug for 5 or 6 days, stop using this medicine, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Precautions for storage and handling

- Store this in a cool place, avoiding direct sunlight.
- Store the product beyond the reach of children.
- After opening, fold up the opening of the bag, and use it as soon as possible.
- Do not transfer the drug to other containers to prevent misuse and quality deterioration.
- Do not use the product past the expiration date.

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.