

## TOKUHON Aroma R

### Indication

stiff shoulders, muscular fatigue, muscular pain, lumbago, contusions, sprains, joint pain, frostbite, bone fracture pain

### Dosage and administration

Peel the film on the drug surface, and adhere it onto the lesion several times daily.

#### Precaution

- The use of the drug for children should be supervised by a parent. Children under 7 years old should not use this drug because it may cause tingling sensations.
- Clean the affected site before using the drug.
- If your skin is sensitive, do not apply the drug to the same site continuously.
- Put a small piece of the drug on the inner arm before use, and confirm that no rash/redness, itching, and eruption occurred.

### Ingredient

In 100g (0.625m<sup>2</sup>) of ointment

Glycol salicylate 5.0g

I-Menthol 1.5g

Vitamin E acetate 2.0g

Glycyrrhetic Acid 0.18g

#### Excipients

Styrene-isoprene-styrene block copolymer, polyisobutylene, liquid paraffin, ester gum, butylated hydroxytoluene, flavoring, phenylethyl alcohol, other one-component

### Precautions

#### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- Do not use in the following sites.
  - Periorbital area, mucosa, etc.
  - Eczema, contact dermatitis, open wounds.

#### Consultation

- The following persons should contact a physician, pharmacist, or registered sales person for a consultation before use.
  - Patients who have experienced allergic symptoms associated with drugs, etc.
- If you have symptoms such as rash/redness, itching, and eruption in or after use, these may be adverse reactions, so immediately discontinue the use, and show this bag to your physician, pharmacist, or registered sales person for a consultation.
- If symptoms do not improve even after using this medicine for 5 to 6 days, stop taking this medicine and consult a physician, pharmacist or registered sales person, being sure to take this bag with you.

### Precautions for storage and handling

- Store this in a cool place, avoiding direct sunlight.
- Store the product beyond the reach of children.
- Once opening, store the unused portion in its original bag and tightly seal the bag by folding the opening over (to maintain product quality).
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)
- Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened.

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.