

# PABRON NUTRITIONAL MEDICINE

## ORAL LIQUID

Newly designated quasi-  
drugs

Health drinks

### Indication

- ☆Feverish wearing disease, during and after illness (or when physical strength is impaired after illness),Loss of appetite, nutritional disorders,Nutritional supplements during physical fatigue,For nutritional supplementation during pregnancy/nursing
- ☆ Nutrients/tonics
- ☆ Weak constitution

### Dosage and administration

Adults (15 years or over): Take 1 bottle (50 mL) per dose once daily.  
Under 15 years: Do not take

### [Precaution]

Comply with the prescribed dosage and administration instructions.  
(Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug.)  
The drug contains a herbal extract. Sedimentation may occur in rare cases. This does not affect drug efficacy.Shake well before taking the drug.

### ingredient and amount

In 1 bottle (50 mL)  
Ginger extract 44.5mg  
Cinnamon Bark Fluid Extract 0.15mL  
Peony Root extract 30mg  
Jujube Extract 80mg  
Glycyrrhiza Extract 25mg  
Taurine 500mg  
Riboflavin sodium phosphate(vitamin B2) 5mg  
Pyridoxine hydrochloride (vitamin B6) 5mg  
Nicotinamide 20mg

### Excipients

Sugar,Polyoxyethylene hydrogenated castor oil,Povidone ,Sodium benzoate,Butyl parahydroxybenzoate,pH adjusting agent,Caramel,flavor

### [Precaution]

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

### Precautions

#### Consultation

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.
  - Skin : rash
  - Gastrointestinal system : feeling of discomfort in the stomach
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.
  - Diarrhea
- When symptoms do not improve even after taking the medicine for a while, stop taking this medicine, and show this product to a physician, pharmacist, or registered salesperson for consultation.

### Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.  
Store the product beyond the reach of children.  
Do not take the product past the expiration date.

**【Disclaimer on Multilingual OTC Product Information】**

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.