

◆PABRON MEDICAL N is a compound of 6 active ingredients including pseudoephedrine hydrochloride and d-chlorpheniramine maleate. A cold medication that is effective on symptoms associated with the common cold, such as runny nose and stuffy nose.

Indication

To alleviate cold symptoms (sore throat, fever, cough, sputum, runny nose, stuffy nose, sneezing, chill (chill due to fever), headache, joint pain, muscle pain)

Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

15 years or over: 2 tablets per dose, 3 times daily

Under 15 years: Do not take

Comply with the prescribed dosage and administration instructions.

•As shown in the figure, press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets.

How to take out the tablets

• (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 2 tablets

Pseudoephedrine hydrochloride 45mg

d-Chlorpheniramine Maleate 1.17mg

Dipotassium Glycyrrhizinate 13.3mg

Ibuprofen 150mg

L-carbocysteine 250mg

Dihydrocodeine Phosphate 8mg

Excipients

cellulose,anhydrous silicic acid ,Hypromellose,Hydroxypropylcellulose,Talc,calcium hydrogen phosphate,Sodium starch glycolate,Mentha oil ,Magnesium stearate

Precautions

When not to use the product

This product should not be used in the following persons:

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

Persons who have had an allergic symptom to this drug or its ingredients.

Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.

Children under 15 years old

Pregnant women who are expected to give birth within 12 weeks.

Persons with the following symptoms:

Urination difficulty due to prostatic hypertrophy

Persons diagnosed as having the following

Hypertension

heart disease

thyroid disease

Diabetes

This drug should not be taken together with the following drugs:

Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines

(internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.) and gastrointestinal analgesics

For those who are lactating, do not use this medicine or avoid lactating while using this medicine.

Do not drink alcohol before and after taking this drug.

● Do not take this medicine for more than 5 days

●After taking this drug, do not drive a car or operate machinery (sleepiness may occur).

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons who have had insomnia, dizziness, feeling of weakness, shivering, or palpitation due to cold medicine, cough medicine, or internal medicine for rhinitis.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

Liver disease

kidney disease

Glaucoma

Systemic lupus erythematosus

Mixed connective tissue disease

Respiratory functional disorder, obstructive sleep apnea syndrome, adiposity

Patients who experienced any of the following diseases.

Gastric/duodenal ulcer

Ulcerative colitis

Crohn's disease

Patients who are being treated with a monoamine oxidase inhibitor (selegiline hydrochloride, etc.)

(Selegiline hydrochloride is used to treat Parkinson's disease.)

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin: rash/redness, itching, and bruises

Gastrointestinal system: nausea/vomiting, loss of appetite, gastric discomfort, stomach pain, stomatitis, heartburn, heaviness in the stomach, gastrointestinal bleeding, abdominal pain, diarrhea, and bloody stool

Neuropsychiatric system

dizziness

Insomnia

Hypersensitivity

Convulsions

Cardiovascular system: palpitations

Respiratory system:

shortness of breath

Urinary system:

urination difficulty

Other: blurred vision, tinnitus, oedema, nosebleeds, bleeding from the gums, defects in hemostasis, bleeding, back pain, excessive drop of body temperature, and general malaise

The following serious symptoms...

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened (to assure the quality).

21

【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.