

# TOKUHON

Class 3 OTC drug

Pain relief/antipruritic/astringent/anti-inflammatory (including cataplasm)

Tokuhon is a plaster containing methyl salicylate that suppresses inflammation and relieves pain, vitamin E that improves blood circulation, and glycyrrhetic acid that alleviates skin inflammation, gentle for skin, and suitable for relieving muscle inflammation and pain. The four corners are rounded to prevent from being peeled off during use.

## Indication

Muscular fatigue  
joint pain, lumbago, muscular pain, pain due to stiff shoulders, contusion pain, bone fracture pain, pain associated with sprain (sprain pain), frostbite,

## Dosage and administration

Peel the paper liner on the drug surface, and adhere it onto the lesion several times daily.  
The use of the drug for children should be supervised by a parent. Children under 7 years old should not use this drug because it may cause tingling sensations.  
Clean the affected site before using the drug.  
If your skin is sensitive, do not apply the drug to the same site continuously.  
Put a small piece of the drug on the inner arm before use, and confirm that no rash/redness, itching, and eruption occurred.

## ingredient and amount

Active ingredients in 100 g (0.54 m<sup>2</sup>) of ointment  
l-Menthol 6.5g  
dl-Camphor 1.1g  
Methyl Salicylate 7.3g  
Tocopherol acetate 1.35g  
Glycyrrhetic Acid 0.18g

## Excipients

Zinc oxide, Butylated hydroxytoluene, Raw rubber, Polyisoprene, Ester gum, Other two-component

## Precautions

### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- The medicine should not be used on the following parts:  
Periorbital area, mucosa, etc.  
Eczema, contact dermatitis, open wounds.

### Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.  
Patients who have experienced allergic symptoms associated with drugs, etc.  
If you have symptoms such as rash/redness, itching, and eruption after use, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered sales person for a consultation.
- If symptoms do not improve even after using this medicine for 5 to 6 days, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

## Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.  
Store the product beyond the reach of children.  
After opening, use it as soon as possible.  
To prevent misuse and quality deterioration, do not transfer the drug to other containers.  
Do not use the product past the expiration date.

**【Disclaimer on Multilingual OTC Product Information】**

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.